

Table 5.15
Cancer of the Cervix Uteri (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2009
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <34 ^e	>=34 ^g	Complete ^h
<u>Race</u>										
All Races ^b	37,983	35,018	36,166	32,413	25,994	20,081	135,337	202,962	44,749	247,711
White ^b	29,788	28,023	29,009	27,086	21,874	16,439	108,627	164,735	37,738	202,473
Black ^b	5,475	4,723	4,389	3,495	2,827	2,717	17,446	25,791	+	+
Asian/Pacific Islander ^c	1,997	1,688	1,758	+	+	+	6,454	+	+	+
Hispanic ^d	8,040	7,423	6,068	+	+	+	25,845	+	+	+

Estimated prevalence percent^a on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f All Ages	
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+		
<u>Race</u>												
All Races ^c	0.0778%	-	0.0002%	0.0088%	0.0794%	0.1569%	0.1623%	0.1393%	0.1241%	0.0734%	0.0755%	
White ^c	0.0791%	-	-	0.0098%	0.0898%	0.1684%	0.1601%	0.1292%	0.1128%	0.0585%	0.0769%	
Black ^c	0.0679%	-	-	0.0088%	0.0599%	0.1261%	0.1624%	0.1506%	0.1418%	0.1364%	0.0718%	
Asian/Pacific Islander ^c	0.0743%	-	-	0.0022%	0.0377%	0.1088%	0.1670%	0.1839%	0.1752%	0.1399%	0.0697%	
Hispanic ^d	0.0956%	-	-	0.0098%	0.0894%	0.2266%	0.2911%	0.2977%	0.2722%	0.1891%	0.1266%	

^a US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.