Table 18.3 Myeloma

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program, 1975-2009 With up to Five Joinpoints, 1992-2009 With up to Three Joinpoints, Females by Race/Ethnicity

	JP Tre	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2000-09	2005-09	
SEER 9 Delay-Adjusted Incidence ^a , 1975-2009															
All Races	1975-09	0.6*											0.6*	0.6*	
White	1975-09	0.5*											0.5*	0.5*	
Black	1975-09	0.9*											0.9*	0.9*	
SEER 13 Delay-Adjusted Incidence ^b , 1992-2009															
All Races	1992-09	0.3											0.3	0.3	
White	1992-09	0.1											0.1	0.1	
Black	1992-09	0.4											0.4	0.4	
SEER 9 Observed Incidence ^a , 1975-2009															
All Races	1975-92	1.1*	1992-09	0.0									0.0	0.0	
White	1975-09	0.3*											0.3*	0.3*	
Black	1975-09	0.7*											0.7*	0.7*	
SEER 13 Observed	Incidence ^b .	1992-200	19												
All Races	1992-09	-0.1	<u> </u>										-0.1	-0.1	
White	1992-09	-0.3											-0.3	-0.3	
White NH ^{ef}	1992-09	-0.3											-0.3	-0.3	
Black	1992-09	0.0											0.0	0.0	
Black NH ^{ef}	1992-09	0.0											0.0	0.0	
API ^e	1992-09	0.5											0.5	0.5	
AI/AN ^{eg}	1992-09	1.7											1.7	1.7	
Hispanic ^f	1992-09	-0.3											-0.3	-0.3	
U.S. Cancer Morta	ality ^c , 1975-	2009													
All Races	1975-93	1.5*	1993-02	-0.5*	2002-09	-2.9*							-2.3*	-2.9*	
White	1975-93	1.4*	1993-02	-0.5*	2002-09	-2.8*							-2.3*	-2.8*	
Black	1975-96	2.0*	1996-09	-2.2*									-2.2*	-2.2*	
U.S. Cancer Mortality ^c , 1992-2009															
All Races	1992-01	-0.2	2001-09	-2.7*									-2.4*	-2.7*	
White	1992-02	-0.4	2002-09	-2.8*									-2.3*	-2.8*	
White NH ^{ef}	1992-02	-0.4	2002-09	-2.9*									-2.3*	-2.9*	
Black	1992-00	0.1	2000-09	-2.9*									-2.9*	-2.9*	
Black NH ^{ef}	1992-99	0.5	1999-09	-2.5*									-2.5*	-2.5*	
API ^e	1992-09	-1.0											-1.0	-1.0	
AI/AN ^{eg}	1992-09	-2.3											-2.3	-2.3	
Hispanic ^f	1992-97	6.1	1997-09	-2.5*									-2.5*	-2.5*	

Joinpoint Regression Program Version 3.5, April 2011, National Cancer Institute. (<u>http://surveillance.cancer.gov/joinpoint</u>/). The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130) Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
- Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).
- Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
- The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.
- API Asian/Pacific Islander, AI/AN American Indian/Alaska Native, NH Non-Hispanic
- Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. The Hispanic and Non-Hispanic mortality trends exclude deaths from CT, DC, ME, MD, MN, NH, NY, ND, SC, OK and VT.
- Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- The APC/AAPC is significantly different from zero (p<.05).
- Joinpoint regression line analysis could not be performed on data series.

National Cancer Institute

а

b

С

d

f

g

*