

Table 18.9

Myeloma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

	Year of Diagnosis																					
	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	68.8	70.7	73.0	71.4	75.5	69.5	72.8	73.4	74.7	71.1	74.8	75.3	74.2	73.0	73.2	74.9	77.6	74.7	76.1	73.9	76.5	76.5
2-year	52.6	55.0	58.5	56.5	60.0	56.6	58.8	59.1	59.2	58.3	61.1	62.6	60.2	60.2	60.0	63.2	65.2	66.5	64.1	65.7	65.6	
3-year	40.3	43.2	45.9	44.0	48.0	45.7	46.6	48.3	48.1	45.8	49.0	51.5	48.2	49.9	49.6	53.4	55.8	58.2	56.3	57.1		
4-year	31.7	34.5	36.2	37.8	38.2	35.1	37.8	37.6	40.5	38.2	38.9	39.8	38.6	41.0	43.9	46.6	47.6	50.4	49.2			
5-year	25.5	27.1	27.7	30.2	31.0	27.5	31.1	31.3	33.6	31.8	31.8	34.0	31.1	34.1	37.1	39.8	41.7	44.5				
6-year	20.2	21.3	22.3	24.1	24.8	21.3	26.8	26.2	28.3	27.1	26.4	29.5	26.7	28.8	32.3	34.7	37.1					
7-year	16.4	16.9	17.7	18.8	21.1	18.7	22.2	21.0	23.6	23.8	22.5	23.9	22.9	24.7	28.7	31.2						
8-year	13.6	13.5	15.1	17.0	17.5	15.8	17.9	18.3	20.4	21.4	19.8	21.7	19.4	22.0	25.2							
9-year	11.5	11.4	12.0	15.3	16.1	13.8	16.0	15.5	18.2	19.9	17.1	19.0	17.7	19.6								
10-year	10.1	9.7	10.5	13.6	14.4	11.9	14.5	13.7	17.0	17.3	15.4	17.1	15.6									
11-year	8.8	8.7	9.2	12.4	12.8	10.8	13.4	13.1	15.8	15.7	14.1	16.0										
12-year	7.9	7.7	8.4	11.5	11.1	9.0	11.9	11.5	14.2	14.9	12.9											
13-year	6.7	6.8	7.6	10.8	10.3	8.2	10.6	11.1	13.4	13.3												
14-year	6.1	6.1	7.0	9.7	9.5	7.7	9.9	10.4	12.2													
15-year	5.4	5.7	6.3	9.1	9.1	7.6	9.0	9.9														
16-year	4.9	5.2	5.7	8.7	8.3	7.4	8.2															
17-year	4.6	4.7	5.4	8.7	8.0	6.6																
18-year	4.0	4.4	5.1	8.0	7.4																	
19-year	3.8	3.8	5.0	7.3																		
20-year	3.3	3.5	4.7																			

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).