

Table 18.10
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	68.7	70.8	73.9	71.0	76.6	73.2	70.5	74.5	75.4	71.9	74.5	76.4	76.1	74.4	74.8	75.6	77.5	76.7	76.2	73.6	77.0	75.6
2-year	52.8	55.4	60.0	56.8	62.0	60.1	57.6	59.7	59.9	59.7	61.4	63.8	61.7	62.2	62.1	62.1	65.9	68.1	63.7	66.4	66.8	
3-year	39.6	43.1	46.6	44.9	49.2	48.9	44.9	47.4	49.6	46.0	50.8	54.0	51.4	52.5	51.3	52.2	56.8	60.2	55.7	57.5		
4-year	31.2	34.1	37.2	37.2	39.6	36.5	36.1	37.3	41.8	39.8	41.0	43.6	43.0	43.2	46.4	46.7	48.1	51.3	49.0			
5-year	24.7	26.0	28.6	30.4	32.8	29.1	29.9	30.5	36.4	33.8	32.5	38.1	34.8	35.9	39.8	41.4	42.3	45.5				
6-year	19.4	21.0	23.7	22.9	26.9	22.4	26.8	27.8	31.6	29.6	26.6	33.7	29.9	29.9	34.9	35.8	37.8					
7-year	15.6	17.1	19.4	18.5	22.6	19.5	22.1	23.2	26.1	25.8	22.4	27.6	25.9	25.6	31.2	32.6						
8-year	13.1	13.8	16.3	17.7	19.6	17.6	17.5	20.2	22.2	24.1	20.3	25.1	21.5	23.0	27.3							
9-year	11.1	11.9	13.0	16.1	18.0	15.9	16.0	16.8	20.0	21.4	18.3	22.8	19.1	19.2								
10-year	9.5	9.9	11.4	14.6	16.0	13.5	14.9	14.8	19.0	18.5	16.4	21.1	17.3									
11-year	7.7	9.0	10.0	12.5	14.1	13.0	13.4	13.9	17.8	17.2	14.9	20.3										
12-year	7.1	7.7	9.4	12.3	12.4	11.6	11.4	11.4	15.5	16.6	14.3											
13-year	5.9	6.9	8.6	11.6	11.8	10.7	10.0	10.9	14.9	14.4												
14-year	5.4	6.1	8.0	10.5	11.1	10.4	9.6	9.9	13.2													
15-year	4.7	5.7	7.3	10.5	10.5	10.2	8.4	9.2														
16-year	4.5	5.2	6.8	10.2	10.0	10.2	7.9															
17-year	4.1	4.9	6.5	9.9	9.6	9.1																
18-year	3.6	4.5	6.2	8.8	8.7																	
19-year	3.4	4.2	6.2	7.8																		
20-year	2.7	3.9	6.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).