

Table 18.11
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Survival Time																						
1-year	69.0	70.5	72.2	71.8	74.2	65.6	75.7	72.5	74.1	70.2	75.0	74.1	72.0	71.4	71.4	74.0	77.7	72.4	76.1	74.3	75.8	77.4
2-year	52.4	54.7	56.9	56.1	57.6	52.9	60.2	58.5	58.4	56.9	60.8	61.3	58.5	57.8	57.5	64.6	64.4	64.7	64.5	65.1	63.9	
3-year	41.0	43.4	45.1	43.0	46.7	42.4	48.7	49.1	46.4	45.5	47.1	48.7	44.5	46.8	47.6	54.8	54.7	55.9	57.1	56.6		
4-year	32.2	35.0	35.2	38.4	36.6	33.5	39.8	37.9	39.1	36.5	36.7	35.5	33.6	38.3	41.1	46.5	46.9	49.3	49.5			
5-year	26.3	28.3	26.8	29.9	28.9	25.9	32.5	31.9	30.6	29.8	31.1	29.5	26.9	31.9	34.1	37.9	41.1	43.3				
6-year	21.1	21.5	20.9	25.4	22.4	20.2	26.8	24.7	24.6	24.4	26.1	24.7	23.0	27.6	29.4	33.4	36.2					
7-year	17.2	16.7	16.1	19.1	19.3	17.9	22.4	19.1	20.9	21.6	22.5	20.0	19.6	23.7	25.8	29.4						
8-year	14.2	13.1	13.8	16.2	14.9	14.0	18.5	16.4	18.3	18.6	19.2	18.0	17.1	20.7	22.9							
9-year	12.0	10.9	11.0	14.4	13.9	11.7	16.0	14.3	16.1	18.1	15.7	14.8	16.0	20.2								
10-year	10.7	9.4	9.6	12.6	12.5	10.3	13.9	12.7	14.9	16.0	14.4	12.8	13.8									
11-year	9.9	8.5	8.4	12.3	11.1	8.5	13.4	12.3	13.4	14.2	13.2	11.5										
12-year	8.8	7.6	7.3	10.7	9.5	6.3	12.5	11.5	12.5	13.1	11.1											
13-year	7.5	6.7	6.6	9.9	8.5	5.7	11.2	11.1	11.6	12.0												
14-year	6.9	6.0	6.0	8.9	7.5	5.0	10.2	10.9	11.0													
15-year	6.0	5.7	5.3	7.7	7.3	5.0	9.7	10.6														
16-year	5.3	5.1	4.6	7.2	6.3	4.2	8.6															
17-year	5.1	4.6	4.4	7.2	6.0	4.0																
18-year	4.5	4.2	4.0	7.2	5.9																	
19-year	4.2	3.4	3.7	6.6																		
20-year	3.8	3.2	3.4																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).