

Table 19.25  
Non-Hodgkin LymphomaEstimated United States Cancer Prevalence Counts<sup>a</sup> on January 1, 2009  
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 <sup>e</sup>	0 to <34 <sup>e</sup>	>=34 <sup>g</sup>	Complete <sup>h</sup>	
Race	Sex											
All Races <sup>b</sup>	Both Sexes	193,568	124,952	73,118	40,328	23,808	13,206	425,784	474,455	9,881	484,336	
	Males	101,809	64,597	37,822	20,781	12,282	6,865	221,874	246,926	5,185	252,111	
	Females	91,759	60,355	35,296	19,547	11,526	6,341	203,910	227,529	4,696	232,225	
White <sup>b</sup>	Both Sexes	169,344	111,105	64,658	36,119	21,607	12,076	375,558	419,930	10,218	430,148	
	Males	89,651	57,853	33,336	18,717	11,169	6,255	196,674	219,545	5,745	225,290	
	Females	79,693	53,252	31,322	17,402	10,438	5,821	178,884	200,385	4,473	204,858	
Black <sup>b</sup>	Both Sexes	15,176	8,849	5,327	2,662	1,414	748	31,735	34,493	788	35,281	
	Males	7,256	4,279	2,863	1,262	730	392	15,520	16,913	326	17,239	
	Females	7,920	4,570	2,464	1,400	684	356	16,215	17,580	462	18,042	
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	5,134	3,241	1,818	+	+	+	11,000	+	+	+	
	Males	2,740	1,632	930	+	+	+	5,705	+	+	+	
	Females	2,394	1,609	888	+	+	+	5,295	+	+	+	
Hispanic <sup>d</sup>	Both Sexes	13,379	8,080	4,315	+	+	+	27,683	+	+	+	
	Males	7,003	4,207	2,320	+	+	+	14,463	+	+	+	
	Females	6,376	3,873	1,995	+	+	+	13,220	+	+	+	

Estimated prevalence percent<sup>a</sup> on January 1, 2009, of the SEER 11 population diagnosed in the previous 19 years  
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted <sup>f</sup>	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
Race	Sex											
All Races <sup>c</sup>	Both Sexes	0.1153%	0.0026%	0.0115%	0.0208%	0.0396%	0.0862%	0.1728%	0.3318%	0.5039%	0.5486%	0.1155%
	Males	0.1221%	0.0033%	0.0150%	0.0252%	0.0458%	0.1019%	0.1966%	0.3775%	0.5621%	0.6296%	0.1323%
	Females	0.1087%	0.0018%	0.0077%	0.0162%	0.0332%	0.0705%	0.1499%	0.2904%	0.4576%	0.5035%	0.1010%
White <sup>c</sup>	Both Sexes	0.1277%	0.0027%	0.0117%	0.0212%	0.0409%	0.0901%	0.1830%	0.3570%	0.5473%	0.5864%	0.1232%
	Males	0.1347%	0.0036%	0.0154%	0.0261%	0.0476%	0.1061%	0.2070%	0.4044%	0.6074%	0.6742%	0.1407%
	Females	0.1207%	0.0018%	0.0077%	0.0159%	0.0337%	0.0735%	0.1592%	0.3127%	0.4983%	0.5374%	0.1077%
Black <sup>c</sup>	Both Sexes	0.0713%	0.0024%	0.0110%	0.0205%	0.0422%	0.0831%	0.1485%	0.2386%	0.2937%	0.2691%	0.0840%
	Males	0.0753%	0.0029%	0.0142%	0.0240%	0.0511%	0.0989%	0.1722%	0.2720%	0.2816%	0.3000%	0.0941%
	Females	0.0677%	0.0019%	0.0077%	0.0172%	0.0344%	0.0693%	0.1286%	0.2129%	0.3021%	0.2547%	0.0757%
Asian/ Pacific Islander <sup>c</sup>	Both Sexes	0.0746%	0.0017%	0.0097%	0.0181%	0.0262%	0.0574%	0.1129%	0.1978%	0.3356%	0.3810%	0.0763%
	Males	0.0793%	0.0022%	0.0119%	0.0208%	0.0274%	0.0669%	0.1273%	0.2261%	0.3958%	0.4352%	0.0879%
	Females	0.0702%	-	0.0073%	0.0154%	0.0250%	0.0486%	0.1004%	0.1737%	0.2900%	0.3474%	0.0670%
Hispanic <sup>d</sup>	Both Sexes	0.0547%	0.0023%	0.0088%	0.0160%	0.0299%	0.0652%	0.1331%	0.2453%	0.3831%	0.3971%	0.0874%
	Males	0.0552%	0.0030%	0.0115%	0.0196%	0.0350%	0.0733%	0.1456%	0.2524%	0.3854%	0.4209%	0.0934%
	Females	0.0541%	0.0015%	0.0060%	0.0118%	0.0239%	0.0562%	0.1210%	0.2392%	0.3813%	0.3832%	0.0816%

<sup>a</sup> US 2009 cancer prevalence counts are based on 2009 cancer prevalence proportions from the SEER registries and 1/1/2009 US population estimates based on the average of 2008 and 2009 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

<sup>b c d</sup> Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

<sup>e</sup> Maximum limited-duration prevalence: 34 years for 1975-2009 SEER 9 data; 19 years for 1990-2009 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

<sup>f</sup> Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

<sup>g h i</sup> (g) Cases diagnosed more than 34 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <34 and >=34. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.