

Table 23.7
Cancer of the Prostate (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2005-2009</u>						
Age at Diagnosis/Death:						
All ages	154.8	146.9	236.0	23.6	21.7	53.1
Under 65	59.1	55.4	106.0	1.8	1.5	4.8
65 and over	816.4	779.7	1,134.1	174.2	161.4	386.9
All ages (IARC world std) ^c	106.1	100.7	168.3	10.2	9.3	24.0
<u>Age-Specific Rates, 2005-2009</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	-	-	-	-	-	-
35-39	1.1	0.9	2.7	0.0	-	-
40-44	10.6	8.8	26.5	0.2	0.2	0.6
45-49	45.5	40.0	99.4	0.8	0.7	2.1
50-54	144.1	132.9	270.2	3.2	2.7	8.7
55-59	330.8	312.7	573.7	9.1	7.6	24.4
60-64	575.6	546.4	987.4	21.8	18.7	58.3
65-69	859.1	822.7	1,329.5	44.8	38.9	116.5
70-74	930.0	895.6	1,275.5	86.5	76.1	218.0
75-79	860.5	821.5	1,066.9	153.9	141.2	340.3
80-84	682.6	640.0	864.9	274.2	256.6	581.5
85+	566.7	535.0	839.7	559.8	532.2	1,187.4

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.