


Table 27.18
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2005-2009

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.36	0.02		
High Five States				
Maine	6.02 ^b	0.27	(01)	38.03 ^c
Vermont	5.89 ^b	0.41	(02)	35.03 ^c
Nevada	5.36 ^b	0.22	(03)	22.93 ^c
Rhode Island	5.27	0.29	(04)	20.66 ^c
Delaware	5.10	0.33	(05)	16.96 ^c
Low Five States				
Mississippi	3.61 ^b	0.16	(47)	-17.37 ^c
Alaska	3.45	0.44	(48)	-20.87 ^c
New Mexico	3.39 ^b	0.18	(49)	-22.22 ^c
Utah	3.36 ^b	0.19	(50)	-22.92 ^c
Hawaii	2.74 ^b	0.19	(51)	-37.29 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.81 ^b	0.12	(45)	-12.66	Montana	4.38	0.28	(23)	0.44
Alaska	3.45	0.44	(48)	-20.87 ^c	Nebraska	3.94	0.20	(41)	-9.73
Arizona	3.89 ^b	0.11	(42)	-10.90	Nevada	5.36 ^b	0.22	(03)	22.93 ^c
Arkansas	4.16	0.16	(32)	-4.76	New Hampshire	4.94	0.27	(09)	13.17
California	4.06 ^b	0.05	(37)	-6.97	New Jersey	4.96 ^b	0.10	(08)	13.57
Colorado	3.83 ^b	0.14	(44)	-12.33	New Mexico	3.39 ^b	0.18	(49)	-22.22 ^c
Connecticut	4.92 ^b	0.16	(10)	12.81	New York	4.38	0.06	(24)	0.33
Delaware	5.10	0.33	(05)	16.96 ^c	North Carolina	4.13	0.10	(33)	-5.30
D.C.	4.36	0.39	(26)	-0.20	North Dakota	4.29	0.33	(27)	-1.59
Florida	4.39	0.06	(22)	0.59	Ohio	5.02 ^b	0.09	(07)	15.08 ^c
Georgia	4.03	0.10	(38)	-7.70	Oklahoma	4.06	0.14	(36)	-6.87
Hawaii	2.74 ^b	0.19	(51)	-37.29 ^c	Oregon	4.73	0.15	(13)	8.43
Idaho	4.36	0.25	(25)	-0.12	Pennsylvania	4.71 ^b	0.08	(16)	7.95
Illinois	4.50	0.08	(21)	3.01	Rhode Island	5.27	0.29	(04)	20.66 ^c
Indiana	4.59	0.12	(18)	5.20	South Carolina	3.99	0.13	(40)	-8.68
Iowa	4.55	0.16	(20)	4.30	South Dakota	4.08	0.29	(35)	-6.52
Kansas	3.87	0.16	(43)	-11.40	Tennessee	4.23	0.12	(30)	-3.02
Kentucky	4.57	0.14	(19)	4.63	Texas	3.79 ^b	0.06	(46)	-13.22
Louisiana	4.27	0.14	(29)	-2.23	Utah	3.36 ^b	0.19	(50)	-22.92 ^c
Maine	6.02 ^b	0.27	(01)	38.03 ^c	Vermont	5.89 ^b	0.41	(02)	35.03 ^c
Maryland	4.72	0.13	(15)	8.06	Virginia	4.12	0.11	(34)	-5.55
Massachusetts	5.10 ^b	0.12	(06)	16.85 ^c	Washington	4.73	0.12	(14)	8.37
Michigan	4.82 ^b	0.10	(12)	10.35	West Virginia	4.68	0.20	(17)	7.21
Minnesota	4.02	0.12	(39)	-7.81	Wisconsin	4.83 ^b	0.12	(11)	10.61
Mississippi	3.61 ^b	0.16	(47)	-17.37 ^c	Wyoming	4.29	0.41	(28)	-1.73
Missouri	4.21	0.11	(31)	-3.43					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.