

Table 33.1
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2003-2009
Both Sexes by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	1,765,890	66.7	0.0	226,940	60.8	0.1	130,058	65.8	0.2	8,480	59.0	0.6	206,044	66.7	0.1
Oral Cavity & Pharynx	40,638	67.7	0.3	4,858	48.4	0.8	3,337	69.7	0.9	207	65.0	4.0	3,650	63.5	0.9
Esophagus	16,851	20.2	0.4	2,376	14.9	0.9	870	18.3	1.7	88	18.2	4.8	1,417	18.6	1.3
Stomach	24,345	27.9	0.3	4,804	29.6	0.8	4,918	37.8	0.8	253	23.2	3.0	5,968	28.5	0.7
Colon & Rectum	169,224	65.2	0.1	24,823	58.7	0.4	15,715	69.3	0.4	1,034	62.7	1.8	19,833	64.2	0.4
Liver & Intrahepatic Bile Duct	24,190	19.0	0.3	4,307	14.8	0.8	6,018	25.4	0.7	338	17.4	2.5	6,292	19.6	0.7
Pancreas	42,912	6.3	0.1	6,465	6.2	0.4	3,480	8.2	0.6	253	5.5	1.8	5,133	7.3	0.5
Larynx	13,027	67.2	0.5	2,451	59.6	1.2	490	70.4	2.4	52	45.0	10.5	1,231	64.3	1.7
Lung & Bronchus	221,498	18.8	0.1	29,556	15.4	0.3	14,423	20.2	0.4	961	13.1	1.4	14,444	17.9	0.4
Melanoma of the Skin	86,274	89.3	0.1	444	71.8	2.5	572	76.1	2.1	106	81.4	4.4	2,883	84.2	0.8
Breast	259,110	88.4	0.1	33,295	78.9	0.3	22,576	91.1	0.2	1,208	85.5	1.3	30,960	87.0	0.2
Cervix uteri	16,800	71.5	0.4	3,139	62.6	1.0	1,897	73.8	1.2	141	60.0	5.0	5,079	74.2	0.7
Corpus & Uterus, NOS	52,327	83.2	0.2	5,857	63.3	0.7	4,507	83.7	0.6	292	84.0	2.6	6,509	81.6	0.6
Ovary ^d	28,063	44.2	0.4	2,757	37.6	1.1	2,370	56.5	1.2	165	46.1	4.4	3,827	49.5	1.0
Prostate	280,354	93.8	0.1	50,548	92.0	0.1	16,275	94.3	0.2	893	90.2	1.2	30,744	92.9	0.2
Testis	13,835	95.8	0.2	422	90.6	1.6	502	93.7	1.3	102	94.0	2.7	3,086	92.8	0.5
Urinary Bladder	76,972	79.0	0.2	4,527	67.5	0.8	3,141	79.9	0.8	176	69.4	4.0	4,867	77.7	0.7
Kidney & Renal Pelvis	54,490	73.1	0.2	6,975	74.2	0.6	3,071	72.5	0.9	495	70.2	2.4	8,252	73.3	0.6
Brain & Nervous System	28,232	33.9	0.3	2,250	42.8	1.2	1,601	43.4	1.4	121	48.0	5.3	4,248	46.4	0.9
Thyroid	45,794	96.8	0.1	3,598	96.3	0.4	5,383	96.4	0.3	230	95.0	1.5	8,045	97.1	0.2
Hodgkin Lymphoma	12,206	87.3	0.3	1,733	83.6	1.0	655	87.6	1.5	42	88.8	5.5	2,276	85.7	0.8
Non-Hodgkin Lymphoma	76,462	70.5	0.2	7,254	65.2	0.6	5,471	67.7	0.7	303	67.4	3.0	10,159	67.7	0.5
Myeloma	20,123	47.6	0.5	5,255	49.6	0.9	1,379	48.9	1.8	110	50.5	6.0	2,967	48.0	1.2
Leukemia	49,389	59.6	0.3	4,661	54.6	0.9	3,204	55.3	1.0	229	60.6	3.5	7,797	61.0	0.6
Mesothelioma	3,991	8.8	0.6	216	14.6	3.2	121	11.4	3.2	-	-	-	440	13.0	2.1

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.2
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2003-2009
Males by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	915,778	66.7	0.1	121,406	63.0	0.2	61,553	61.5	0.2	4,063	56.8	0.9	101,466	65.6	0.2
Oral Cavity & Pharynx	28,796	67.7	0.3	3,334	43.5	1.0	2,154	67.5	1.2	141	63.2	5.1	2,459	60.3	1.2
Esophagus	13,327	20.6	0.4	1,642	13.3	1.1	657	18.9	1.9	56	14.6	5.9	1,133	18.4	1.4
Stomach	15,014	26.0	0.4	2,693	27.3	1.0	2,744	37.9	1.1	160	20.1	3.6	3,382	28.4	1.0
Colon & Rectum	87,211	65.8	0.2	11,723	57.7	0.6	8,104	69.2	0.6	518	61.8	2.6	10,680	64.3	0.6
Liver & Intrahepatic Bile Duct	17,563	19.2	0.4	3,159	12.8	0.9	4,217	26.0	0.9	240	20.8	3.1	4,521	19.4	0.8
Pancreas	21,391	5.9	0.2	2,949	5.8	0.6	1,702	7.6	0.8	123	3.7	2.2	2,463	6.2	0.6
Larynx	10,487	67.9	0.5	1,985	59.3	1.3	426	69.9	2.6	41	46.2	12.4	1,059	63.6	1.9
Lung & Bronchus	116,718	16.4	0.1	16,770	13.6	0.3	8,415	17.5	0.5	499	13.9	2.0	7,871	15.0	0.5
Melanoma of the Skin	48,363	87.0	0.2	186	62.1	4.1	282	69.4	3.3	54	69.9	7.9	1,188	78.2	1.5
Prostate	280,354	93.8	0.1	50,548	92.0	0.1	16,275	94.3	0.2	893	90.2	1.2	30,744	92.9	0.2
Testis	13,835	95.8	0.2	422	90.6	1.6	502	93.7	1.3	102	94.0	2.7	3,086	92.8	0.5
Urinary Bladder	58,165	80.4	0.2	2,971	71.7	1.0	2,385	81.5	0.9	140	71.7	4.5	3,584	79.7	0.8
Kidney & Renal Pelvis	33,621	72.7	0.3	4,168	73.4	0.8	1,921	72.2	1.2	291	70.1	3.2	4,886	71.9	0.8
Brain & Nervous System	15,775	33.0	0.4	1,134	38.5	1.7	848	41.4	2.0	59	56.1	6.8	2,325	45.2	1.2
Thyroid	10,759	94.0	0.3	657	93.2	1.2	1,065	94.6	0.8	44	88.1	5.0	1,428	94.0	0.8
Hodgkin Lymphoma	6,637	87.1	0.5	913	80.9	1.5	359	86.2	2.1	-	-	-	1,263	85.0	1.2
Non-Hodgkin Lymphoma	41,129	69.6	0.3	3,919	61.7	0.9	2,917	66.2	1.0	161	68.9	4.3	5,407	65.8	0.7
Myeloma	11,182	49.2	0.6	2,555	48.3	1.3	723	49.8	2.5	52	49.5	8.5	1,557	50.7	1.7
Leukemia	28,377	60.3	0.3	2,521	55.3	1.2	1,783	56.5	1.3	127	62.0	4.8	4,364	62.5	0.8
Mesothelioma	3,054	6.9	0.6	151	12.7	4.0	88	7.6	3.1	-	-	-	304	11.4	2.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.3
Five-Year Cause-Specific Survival^a(%) for Malignant Cancers, 2003-2009
Females by Race/Ethnicity and Primary Site

Primary Site	White			Black			Asian/ Pacific Islander			American Indian/ Alaska Native ^b			Hispanic ^c		
	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE	Cases	Percent	SE
All Sites	850,112	66.7	0.1	105,534	58.3	0.2	68,505	69.5	0.2	4,417	61.0	0.8	104,578	67.8	0.2
Oral Cavity & Pharynx	11,842	67.7	0.5	1,524	58.7	1.5	1,183	73.6	1.5	66	68.6	6.6	1,191	70.2	1.6
Esophagus	3,524	19.0	0.8	734	18.5	1.7	213	16.7	3.3	32	23.6	7.7	284	19.3	2.7
Stomach	9,331	30.9	0.6	2,111	32.7	1.2	2,174	37.7	1.2	93	28.9	5.2	2,586	28.7	1.1
Colon & Rectum	82,013	64.7	0.2	13,100	59.5	0.5	7,611	69.5	0.6	516	63.4	2.4	9,153	64.2	0.6
Liver & Intrahepatic Bile Duct	6,627	18.2	0.6	1,148	19.6	1.5	1,801	24.2	1.3	98	7.3	4.2	1,771	20.0	1.2
Pancreas	21,521	6.7	0.2	3,516	6.6	0.5	1,778	8.9	0.8	130	7.4	2.7	2,670	8.3	0.7
Larynx	2,540	64.3	1.1	466	60.8	2.7	64	74.1	6.3	-	-	-	172	68.4	4.4
Lung & Bronchus	104,780	21.5	0.2	12,786	17.7	0.4	6,008	23.9	0.7	462	12.6	1.9	6,573	21.4	0.6
Melanoma of the Skin	37,911	92.2	0.2	258	78.8	2.9	290	82.8	2.5	52	91.9	3.9	1,695	88.2	0.9
Breast	257,235	88.5	0.1	32,996	78.9	0.3	22,457	91.1	0.2	1,203	85.4	1.3	30,805	87.0	0.2
Cervix uteri	16,800	71.5	0.4	3,139	62.6	1.0	1,897	73.8	1.2	141	60.0	5.0	5,079	74.2	0.7
Corpus & Uterus, NOS	52,327	83.2	0.2	5,857	63.3	0.7	4,507	83.7	0.6	292	84.0	2.6	6,509	81.6	0.6
Ovary ^d	28,063	44.2	0.4	2,757	37.6	1.1	2,370	56.5	1.2	165	46.1	4.4	3,827	49.5	1.0
Urinary Bladder	18,807	74.8	0.4	1,556	59.5	1.5	756	75.0	1.8	36	60.3	8.3	1,283	72.2	1.4
Kidney & Renal Pelvis	20,869	73.8	0.3	2,807	75.3	1.0	1,150	73.2	1.5	204	69.9	3.5	3,366	75.4	0.9
Brain & Nervous System	12,457	35.1	0.5	1,116	47.1	1.7	753	45.6	2.1	62	42.0	7.4	1,923	47.9	1.3
Thyroid	35,035	97.7	0.1	2,941	96.9	0.3	4,318	96.9	0.3	186	96.6	1.4	6,617	97.8	0.2
Hodgkin Lymphoma	5,569	87.7	0.5	820	86.5	1.4	296	89.4	1.9	-	-	-	1,013	86.6	1.2
Non-Hodgkin Lymphoma	35,333	71.4	0.3	3,335	69.3	0.9	2,554	69.4	1.0	142	65.4	4.3	4,752	69.9	0.8
Myeloma	8,941	45.6	0.7	2,700	50.8	1.3	656	48.0	2.6	58	51.3	8.3	1,410	45.1	1.8
Leukemia	21,012	58.7	0.4	2,140	53.7	1.3	1,421	53.9	1.5	102	59.1	5.2	3,433	59.0	1.0
Mesothelioma	937	15.0	1.5	65	16.9	6.0	33	21.8	7.8	-	-	-	136	17.0	4.1

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^c Hispanics are not mutually exclusive from whites, blacks, Asian/Pacific Islanders and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

- Statistic could not be calculated due to less than 25 cases in the time interval.

Table 33.4

Cancer of the Female Breast

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2003-2009

Race/Ethnicity	All Ages				Ages <50				Ages 50+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	316,365	100.0	87.7	0.1	79,603	100.0	89.4	0.1	236,762	100.0	87.1	0.1
White	257,235	81.3	88.5	0.1	60,495	76.0	90.6	0.1	196,740	83.1	87.8	0.1
White Non-Hispanic	227,585	71.9	88.6	0.1	49,511	62.2	91.2	0.2	178,074	75.2	87.9	0.1
White Hispanic	29,650	9.4	87.1	0.2	10,984	13.8	87.7	0.4	18,666	7.9	86.7	0.3
Black	32,996	10.4	78.9	0.3	10,367	13.0	79.7	0.5	22,629	9.6	78.6	0.3
American Indian/Alaska Native ^c	1,203	0.4	85.4	1.3	386	0.5	86.4	2.0	817	0.3	84.9	1.6
Asian	20,787	6.6	91.4	0.2	7,120	8.9	92.9	0.4	13,667	5.8	90.7	0.3
Asian Indian, Pakistani	1,696	0.5	89.7	1.0	668	0.8	92.6	1.3	1,028	0.4	87.7	1.3
Chinese	4,147	1.3	91.7	0.5	1,504	1.9	93.3	0.8	2,643	1.1	90.9	0.7
Filipino	5,721	1.8	90.0	0.5	1,713	2.2	90.6	0.9	4,008	1.7	89.7	0.6
Japanese	3,148	1.0	92.8	0.5	701	0.9	95.8	1.0	2,447	1.0	92.0	0.6
Korean	1,532	0.5	92.1	0.8	673	0.8	93.7	1.1	859	0.4	90.8	1.2
Vietnamese	1,337	0.4	91.4	1.0	570	0.7	92.8	1.4	767	0.3	90.4	1.4
Other Asian ^d	3,206	1.0	92.8	0.6	1,291	1.6	93.8	0.8	1,915	0.8	92.1	0.7
Pacific Islander	1,670	0.5	86.8	1.1	504	0.6	83.6	2.1	1,166	0.5	88.3	1.2
Hawaiian	1,087	0.3	89.7	1.1	302	0.4	88.4	2.3	785	0.3	90.3	1.3
Other Pacific Islander ^e	583	0.2	80.8	2.2	202	0.3	75.6	4.0	381	0.2	83.6	2.5
Hispanic ^f	30,805	9.7	87.0	0.2	11,392	14.3	87.7	0.4	19,413	8.2	86.6	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.5

Cancer of the Colon and Rectum

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2003-2009

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	212,664	100.0	64.9	0.1	108,602	100.0	65.4	0.2	104,062	100.0	64.5	0.2
White	169,224	79.6	65.2	0.1	87,211	80.3	65.8	0.2	82,013	78.8	64.7	0.2
White Non-Hispanic	150,136	70.6	65.4	0.1	76,894	70.8	66.0	0.2	73,242	70.4	64.7	0.2
White Hispanic	19,088	9.0	64.2	0.4	10,317	9.5	64.1	0.6	8,771	8.4	64.3	0.6
Black	24,823	11.7	58.7	0.4	11,723	10.8	57.7	0.6	13,100	12.6	59.5	0.5
American Indian/Alaska Native ^c	1,034	0.5	62.7	1.8	518	0.5	61.8	2.6	516	0.5	63.4	2.4
Asian	14,797	7.0	69.9	0.5	7,563	7.0	70.0	0.6	7,234	7.0	69.8	0.6
Asian Indian, Pakistani	696	0.3	73.4	2.0	410	0.4	73.8	2.6	286	0.3	73.0	3.1
Chinese	3,463	1.6	70.9	0.9	1,736	1.6	71.6	1.3	1,727	1.7	70.3	1.3
Filipino	3,190	1.5	67.8	1.0	1,651	1.5	68.1	1.4	1,539	1.5	67.5	1.5
Japanese	2,942	1.4	69.3	1.0	1,473	1.4	68.2	1.4	1,469	1.4	70.3	1.4
Korean	1,440	0.7	69.5	1.4	709	0.7	69.4	2.1	731	0.7	69.7	2.0
Vietnamese	1,205	0.6	68.5	1.6	641	0.6	67.8	2.4	564	0.5	69.1	2.3
Other Asian ^d	1,861	0.9	72.5	1.2	943	0.9	73.6	1.7	918	0.9	71.3	1.8
Pacific Islander	918	0.4	59.8	2.0	541	0.5	57.6	2.7	377	0.4	63.0	3.1
Hawaiian	624	0.3	59.9	2.5	374	0.3	56.2	3.3	250	0.2	65.6	3.8
Other Pacific Islander ^e	294	0.1	59.5	3.6	167	0.2	60.9	4.7	127	0.1	57.9	5.3
Hispanic ^f	19,833	9.3	64.2	0.4	10,680	9.8	64.3	0.6	9,153	8.8	64.2	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.6

Cancer of the Liver and Intrahepatic Bile Duct

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2003-2009

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	35,081	100.0	19.7	0.3	25,349	100.0	19.7	0.3	9,732	100.0	19.5	0.5
White	24,190	69.0	19.0	0.3	17,563	69.3	19.2	0.4	6,627	68.1	18.2	0.6
White Non-Hispanic	18,091	51.6	18.8	0.4	13,170	52.0	19.2	0.4	4,921	50.6	17.9	0.7
White Hispanic	6,099	17.4	19.4	0.7	4,393	17.3	19.4	0.8	1,706	17.5	19.3	1.2
Black	4,307	12.3	14.8	0.8	3,159	12.5	12.8	0.9	1,148	11.8	19.6	1.5
American Indian/Alaska Native ^c	338	1.0	17.4	2.5	240	0.9	20.8	3.1	98	1.0	7.3	4.2
Asian	5,729	16.3	25.9	0.7	3,994	15.8	26.6	0.9	1,735	17.8	24.5	1.3
Asian Indian, Pakistani	179	0.5	23.2	4.2	122	0.5	24.7	5.4	57	0.6	18.7	6.7
Chinese	1,469	4.2	28.4	1.5	1,043	4.1	29.3	1.8	426	4.4	26.1	2.9
Filipino	859	2.4	23.1	1.8	617	2.4	23.0	2.2	242	2.5	23.2	3.3
Japanese	552	1.6	23.8	2.3	269	1.1	30.3	3.4	283	2.9	19.6	3.0
Korean	684	1.9	27.3	2.0	458	1.8	25.3	2.4	226	2.3	31.5	3.6
Vietnamese	1,221	3.5	28.1	1.6	941	3.7	28.1	1.8	280	2.9	27.7	3.5
Other Asian ^d	765	2.2	21.8	1.9	544	2.1	23.1	2.2	221	2.3	17.6	3.8
Pacific Islander	289	0.8	15.2	2.8	223	0.9	14.9	3.1	66	0.7	16.1	6.4
Hawaiian	155	0.4	13.2	3.5	116	0.5	14.3	4.1	39	0.4	8.8	7.0
Other Pacific Islander ^e	134	0.4	18.9	4.7	107	0.4	16.3	5.1	27	0.3	33.7	10.0
Hispanic ^f	6,292	17.9	19.6	0.7	4,521	17.8	19.4	0.8	1,771	18.2	20.0	1.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.7

Cancer of the Lung and Bronchus

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2003-2009

Race/Ethnicity	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	267,363	100.0	18.5	0.1	142,890	100.0	16.2	0.1	124,473	100.0	21.2	0.1
White	221,498	82.8	18.8	0.1	116,718	81.7	16.4	0.1	104,780	84.2	21.5	0.2
White Non-Hispanic	207,570	77.6	18.9	0.1	109,118	76.4	16.5	0.1	98,452	79.1	21.5	0.2
White Hispanic	13,928	5.2	17.9	0.4	7,600	5.3	15.0	0.5	6,328	5.1	21.4	0.6
Black	29,556	11.1	15.4	0.3	16,770	11.7	13.6	0.3	12,786	10.3	17.7	0.4
American Indian/Alaska Native ^c	961	0.4	13.1	1.4	499	0.3	13.9	2.0	462	0.4	12.6	1.9
Asian	13,200	4.9	20.4	0.5	7,696	5.4	17.7	0.6	5,504	4.4	24.3	0.8
Asian Indian, Pakistani	528	0.2	26.4	2.5	355	0.2	22.4	2.8	173	0.1	35.2	4.8
Chinese	3,356	1.3	20.5	0.9	1,938	1.4	17.5	1.1	1,418	1.1	24.8	1.5
Filipino	3,393	1.3	21.1	0.9	2,052	1.4	17.8	1.1	1,341	1.1	26.3	1.6
Japanese	2,133	0.8	18.0	1.0	1,110	0.8	16.3	1.4	1,023	0.8	19.8	1.6
Korean	1,047	0.4	18.4	1.5	585	0.4	16.2	1.9	462	0.4	20.9	2.4
Vietnamese	1,476	0.6	20.0	1.4	959	0.7	16.9	1.5	517	0.4	26.0	2.6
Other Asian ^d	1,267	0.5	22.3	1.5	697	0.5	20.1	2.0	570	0.5	25.0	2.3
Pacific Islander	1,223	0.5	17.6	1.4	719	0.5	15.9	1.7	504	0.4	20.0	2.2
Hawaiian	843	0.3	17.5	1.6	478	0.3	15.7	2.1	365	0.3	19.7	2.5
Other Pacific Islander ^e	380	0.1	17.7	2.6	241	0.2	16.2	3.1	139	0.1	20.5	4.5
Hispanic ^f	14,444	5.4	17.9	0.4	7,871	5.5	15.0	0.5	6,573	5.3	21.4	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorroan, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.

Table 33.8

Cancer of the Prostate

5-Year Cause-Specific Survival^a and Case Distribution by Race/Ethnicity
SEER 18 Registries, 2003-2009

Race/Ethnicity	All Ages				Ages <65				Ages 65+			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE	Count	Percent ^b	Percent	SE
All Races	359,223	100.0	93.7	0.0	152,715	100.0	96.6	0.1	206,508	100.0	91.5	0.1
White	280,354	78.0	93.8	0.1	116,610	76.4	96.9	0.1	163,744	79.3	91.6	0.1
White Non-Hispanic	251,360	70.0	93.9	0.1	104,727	68.6	97.1	0.1	146,633	71.0	91.7	0.1
White Hispanic	28,994	8.1	92.9	0.2	11,883	7.8	95.5	0.2	17,111	8.3	91.1	0.3
Black	50,548	14.1	92.0	0.1	26,576	17.4	95.1	0.2	23,972	11.6	88.5	0.3
American Indian/Alaska Native ^c	893	0.2	90.2	1.2	382	0.3	94.0	1.4	511	0.2	87.2	1.8
Asian	15,137	4.2	94.6	0.2	4,836	3.2	97.0	0.3	10,301	5.0	93.5	0.3
Asian Indian, Pakistani	1,381	0.4	96.0	0.6	568	0.4	97.9	0.7	813	0.4	94.6	1.0
Chinese	3,271	0.9	95.0	0.5	889	0.6	98.8	0.5	2,382	1.2	93.7	0.6
Filipino	4,238	1.2	93.4	0.5	1,468	1.0	95.0	0.8	2,770	1.3	92.6	0.6
Japanese	2,693	0.7	94.8	0.5	667	0.4	97.3	0.8	2,026	1.0	93.9	0.6
Korean	736	0.2	94.9	1.0	182	0.1	95.5	2.0	554	0.3	94.7	1.1
Vietnamese	828	0.2	94.2	1.0	290	0.2	96.6	1.2	538	0.3	92.9	1.3
Other Asian ^d	1,990	0.6	95.4	0.6	772	0.5	98.3	0.6	1,218	0.6	93.6	0.8
Pacific Islander	1,138	0.3	90.2	1.1	404	0.3	93.0	1.5	734	0.4	88.7	1.4
Hawaiian	672	0.2	89.8	1.4	238	0.2	92.1	2.0	434	0.2	88.7	1.9
Other Pacific Islander ^e	466	0.1	90.9	1.7	166	0.1	94.4	2.1	300	0.1	88.8	2.3
Hispanic ^f	30,744	8.6	92.9	0.2	12,495	8.2	95.5	0.2	18,249	8.8	91.2	0.3

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^a Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

^b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

^c Survival estimates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

^e Other Pacific Islanders include Micronesian, Chamorro, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

^f Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic not shown due to fewer than 25 cases during the time period.