

Table 1.23
U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2010^a
By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
All Sites									
Males	6,078,974	18,390	42,087	83,873	155,608	354,801	860,143	1,578,142	2,985,931
Females	6,948,940	16,117	36,351	86,197	226,859	636,726	1,276,932	1,684,808	2,984,950
Oral Cavity & Pharynx									
Males	181,084	68	468	1,480	3,518	14,224	43,847	54,509	62,971
Females	94,109	90	553	1,470	3,635	8,681	17,691	22,821	39,167
Esophagus									
Males	26,388	0	0	22	219	1,177	4,653	9,419	10,897
Females	7,451	0	0	0	28	256	1,263	1,790	4,114
Stomach									
Males	41,386	0	18	64	607	2,589	7,095	10,242	20,773
Females	30,883	4	12	165	568	2,307	4,398	6,312	17,118
Colon & Rectum									
Males	569,930	11	117	1,190	5,632	25,029	80,675	136,002	321,273
Females	584,551	0	75	1,251	5,467	23,433	67,578	110,244	376,504
Liver & Intrahep									
Males	28,967	484	535	554	454	1,582	10,000	9,083	6,274
Females	12,437	471	470	312	411	895	2,803	3,028	4,048
Pancreas									
Males	20,293	11	12	127	376	1,616	4,320	6,183	7,649
Females	21,316	0	66	155	413	1,376	3,843	5,483	9,980
Larynx									
Males	71,659	0	0	78	243	1,915	10,556	20,739	38,129
Females	17,370	0	0	25	134	1,011	3,256	4,677	8,267
Lung & Bronchus									
Males	183,410	45	90	444	1,184	5,545	24,951	54,888	96,262
Females	216,021	31	71	396	1,435	7,883	28,898	59,047	118,261
Melanoma of the Skin									
Males	452,540	78	656	5,014	18,294	46,766	95,157	122,709	163,865
Females	469,240	113	993	10,894	33,972	73,514	109,961	106,450	133,343

^a U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table 1.23 - continued
 U.S. Complete Prevalence Counts, Invasive Cancers Only, January 1, 2010^a
 By Age at Prevalence

Site/Sex	Age at Prevalence								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
Breast									
Males	14,588	0	0	11	89	586	2,018	4,133	7,751
Females	2,829,041	0	53	2,740	36,291	216,912	539,229	767,846	1,265,969
Cervix									
Females	249,496	0	45	2,239	16,405	42,748	59,766	55,248	73,044
Corpus & Uterus, NOS									
Females	600,346	0	47	530	5,666	24,621	87,167	157,833	324,482
Ovary ^d									
Females	186,138	84	1,018	3,453	6,883	18,736	40,700	47,632	67,632
Prostate									
Males	2,617,682	34	59	101	324	21,033	228,561	740,428	1,627,144
Urinary Bladder									
Males	419,648	45	105	611	2,438	11,535	42,898	99,644	262,372
Females	143,992	23	57	217	964	4,266	13,698	30,030	94,737
Kidney & Renal Pelvis									
Males	202,516	1,454	2,505	2,659	5,172	17,901	40,060	57,511	75,253
Females	138,989	1,343	2,681	2,851	4,901	12,071	23,973	33,716	57,452
Hodgkin Lymphoma									
Males	93,890	181	2,315	9,272	16,487	24,007	21,438	13,274	6,916
Females	88,038	101	1,901	9,617	16,530	23,444	18,638	10,651	7,157
Non-Hodgkin Lymphoma									
Males	266,478	773	3,513	7,582	13,366	29,224	52,496	67,493	92,032
Females	242,587	430	1,847	4,348	8,879	20,902	41,030	56,723	108,427
Myeloma									
Males	42,185	0	6	84	442	2,859	8,141	13,608	17,045
Females	35,432	0	0	39	281	2,266	6,273	10,676	15,896
Leukemia									
Males	162,651	6,545	12,947	12,388	10,928	13,500	22,064	32,827	51,452
Females	125,312	5,438	10,680	10,750	9,514	9,728	14,680	20,623	43,899
Acute Lymphocytic Leuk									
Males	38,239	5,461	11,081	9,597	6,075	3,568	1,238	775	444
Females	31,328	4,527	9,014	7,859	5,163	2,755	884	717	408

^a U.S. 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2010 U.S. population estimates based on the average of 2009 and 2010 population estimates from the U.S. Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.