- - - -

Table 1.25

Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity

All Races			White			Black		
	Rate ^b 2006-2010	APC ^c 2001-2010		Rate ^b 2006-2010	APC ^c 2001-2010		Rate ^b 2006-2010	APC ^c 2001-2010
All Sites	535.9	-1.1*	All Sites	539.1	-1.1*	All Sites	610.4	-1.9*
Prostate	152.0	-2.3*	Prostate	144.9	-2.5*	Prostate	228.5	-2.6*
Lung and Bronchus	74.3	-2.2*	Lung and Bronchus	74.5	-2.2*	Lung and Bronchus	95.8	-2.8*
Colon and Rectum	52.2	-2.9*	Colon and Rectum	51.3	-3.1*	Colon and Rectum	64.3	-2.4*
Urinary Bladder	36.6	-0.7*	Urinary Bladder	39.8	-0.7*	Kidney and Renal Pelvis	24.1	2.5*
Melanoma of the Skin	27.4	1.8*	Melanoma of the Skin	31.9	1.8*	Urinary Bladder	21.0	0.4
Non-Hodgkin Lymphoma	23.9	0.2	Non-Hodgkin Lymphoma	25.0	0.2	Pancreas	17.6	1.3*
Kidney and Renal Pelvis	21.0	2.2*	Kidney and Renal Pelvis	21.5	2.2*	Non-Hodgkin Lymphoma	17.4	-0.2
Leukemia	16.3	-0.8*	Leukemia	17.2	-0.8*	Stomach	16.1	-2.1*
Oral Cavity and Pharynx	16.2	0.2	Oral Cavity and Pharynx	16.7	0.6*	Liver & IBD ^f	15.1	3.7*
Pancreas	13.9	0.8*	Pancreas	13.8	0.8*	Oral Cavity and Pharynx	14.9	-3.1*
Liver & IBD ^f	11.9	3.8*	Liver & IBD ^f	10.4	4.5*	Myeloma	14.4	-0.4
Stomach	10.4	-1.5*	Stomach	9.2	-1.3*	Leukemia	12.6	-1.9*
Esophagus	7.7	-0.9*	Brain and ONS ^f	8.4	-0.1	Larynx	9.6	-3.7*
Brain and ONS ^f	7.7	-0.3	Esophagus	8.0	-0.2	Esophagus	8.4	-5.0*
Myeloma	7.5	0.6*	Myeloma	7.1	0.7*	Brain and ONS ^f	4.7	-0.2
Asian/Pacific Islander			American Indian/Alaska Native ^d			Hispanic ^e		
	Rateb	APC°	·	Rateb	APC°		Rateb	APC°
	2006-2010	2001-2010		2006-2010			2006-2010	2001-2010
All Sites	335.0	-1.8*	All Sites	351.3	-0.7	All Sites	409.7	-1.4*
Prostate	81.8	-3.8*	Prostate	77.8	-2.2*	Prostate	125.8	-3.0*
Lung and Bronchus	50.7	-1.7*	Lung and Bronchus	51.2	-2.3*	Colon and Rectum	45.5	-1.7*
Colon and Rectum	43.8	-2.7*	Colon and Rectum	44.1	-1.2	Lung and Bronchus	40.6	-2.6*
Liver & IBD ^f	21.4	-0.9	Kidney and Renal Pelvis	24.8	2.9	Non-Hodgkin Lymphoma	20.5	0.7
Urinary Bladder	16.1	-1.0*	Liver & IBDf	20.6	6.1	Urinary Bladder	20.2	-1.5*
Non-Hodgkin Lymphoma	16.1	-0.2	Non-Hodgkin Lymphoma	14.1	-1.1	Kidney and Renal Pelvis	20.1	2.1*
Stomach	15.5	-3.5*	Urinary Bladder	13.7	_	Liver & IBDf	18.3	3.2*
Kidney and Renal Pelvis	11.0	3.2*	Stomach	13.1	-5.4*	Stomach	14.9	-1.3*
Oral Cavity and Pharynx	11.0	0.4	Pancreas	11.4	_	Leukemia	12.2	-0.3
Pancreas	10.4	0.3	Oral Cavity and Pharynx	11.1	2.9	Pancreas	12.1	0.9
Leukemia	9.0	-0.8	Leukemia	8.8	_	Oral Cavity and Pharynx	9.3	-0.1
Thyroid	5.3	4.6*	Esophagus	6.1	_	Myeloma	6.7	-1.3
Myeloma	4.3	1.5	Larynx	4.6	_	Brain and ONS ^f	5.9	-1.3*
Brain and ONSf	4.2	0.1	Testis	4.5	_	Esophagus	5.2	-1.0
Esophagus	3.9	-1.3	Melanoma of the Skin	4.1	-	Melanoma of the Skin	4.7	-1.2

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

- a Top 15 cancer sites selected based on 2006-2010 age-adjusted rates for the race/ethnic group.
- Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.