

Table 2.5  
All Cancer Sites (Invasive)

Age-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Year of Diagnosis:									
1975-2010	464.56	553.17	407.65	470.43	555.86	417.20	508.16	672.16	399.24
1975	400.44	466.85	365.86	402.12	468.61	369.55	426.53	525.51	356.64
1976	407.40	481.12	367.43	409.72	481.86	373.16	427.31	552.85	340.41
1977	407.77	486.55	363.72	409.34	487.20	367.90	442.64	577.58	352.06
1978	407.33	488.39	362.23	409.84	489.80	368.00	436.29	570.52	343.84
1979	412.55	496.47	365.61	414.17	497.64	369.77	451.71	589.96	359.29
1980	418.06	505.82	368.10	419.48	505.21	373.36	467.54	615.22	366.55
1981	425.30	510.88	377.15	427.73	510.90	383.94	472.74	633.91	362.94
1982	424.51	511.10	376.22	427.84	512.53	383.51	465.14	621.86	359.38
1983	431.18	520.00	380.79	434.42	521.02	387.98	486.95	645.95	381.89
1984	440.06	525.52	392.53	443.73	527.79	399.64	488.15	643.59	386.18
1985	448.95	531.34	403.24	453.56	533.67	411.57	489.03	643.53	388.23
1986	451.44	538.27	402.38	456.07	542.94	409.21	493.75	643.87	396.43
1987	468.05	563.22	413.29	474.88	568.82	423.30	501.00	666.16	394.10
1988	463.81	558.35	409.21	470.26	565.05	417.82	503.73	657.40	404.15
1989	467.79	567.10	408.34	475.17	573.58	418.60	492.46	661.26	384.33
1990	481.97	592.68	415.55	488.88	599.24	424.85	521.51	699.38	408.65
1991	503.03	637.40	418.94	510.49	644.48	428.54	548.71	764.40	412.31
1992	510.56	658.12	414.82	516.76	663.30	423.44	568.46	808.22	413.25
1993	493.37	622.04	408.30	496.89	620.13	417.05	567.86	817.52	403.82
1994	483.46	589.63	413.52	487.33	587.37	423.05	557.61	782.98	408.81
1995	476.90	571.59	414.82	482.09	570.05	426.09	542.24	747.92	404.48
1996	479.13	572.50	418.05	484.73	573.78	428.03	540.00	726.00	416.83
1997	486.35	575.53	428.33	493.45	576.29	441.08	539.99	739.45	409.83
1998	488.19	572.75	433.78	496.76	576.39	446.89	535.30	721.81	415.21
1999	490.38	580.47	430.49	498.38	582.43	443.51	539.56	721.82	420.01
2000	486.04	579.89	421.98	496.27	583.60	437.68	526.20	710.72	402.82
2001	489.68	580.98	426.86	500.65	587.01	442.11	519.31	696.54	399.92
2002	487.48	574.46	427.95	497.43	580.83	440.79	533.43	696.50	424.68
2003	475.18	559.75	417.11	485.69	564.54	432.15	517.90	678.19	411.28
2004	476.14	557.70	419.72	486.11	563.09	433.05	522.60	681.22	418.23
2005	471.86	548.27	418.88	483.55	556.61	433.29	505.56	637.84	418.04
2006	475.02	554.58	418.29	486.24	561.28	432.91	505.08	639.30	413.34
2007	480.55	565.07	419.50	490.79	570.11	433.71	513.30	660.63	412.22
2008	473.43	545.48	422.19	482.94	549.48	435.95	512.31	656.60	414.46
2009	470.46	534.42	425.19	479.12	538.31	437.45	501.62	627.67	416.29
2010	457.54	519.27	413.32	465.44	521.40	425.80	479.73	597.48	398.51

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
- Statistic not shown. Rate based on less than 16 cases for the time interval.