

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2006-2010</u>						
Age at Diagnosis/Death:						
All ages	7.9	7.9	9.6	2.4	2.2	4.2
Under 65	7.4	7.5	7.9	1.8	1.7	2.8
65 and over	11.7	10.5	21.0	6.0	5.2	13.3
All ages (IARC world std) ^c	6.3	6.3	7.3	1.7	1.6	3.0
<u>Age-Specific Rates, 2006-2010</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.1	0.2	-	-	-	-
20-24	1.4	1.4	1.5	0.1	0.1	0.2
25-29	5.1	5.6	4.3	0.5	0.5	0.8
30-34	10.3	11.2	9.2	1.3	1.2	1.8
35-39	14.3	15.6	11.0	2.4	2.4	3.2
40-44	15.1	15.5	16.3	3.2	3.0	4.9
45-49	14.0	13.9	14.3	4.0	3.7	6.2
50-54	13.1	12.5	16.8	4.6	4.2	7.3
55-59	12.0	11.3	17.4	4.8	4.5	7.4
60-64	12.9	11.8	18.8	5.2	4.7	9.1
65-69	13.4	12.4	20.3	5.5	4.8	11.6
70-74	12.8	11.5	19.9	5.7	5.1	11.5
75-79	10.5	9.4	19.4	5.8	5.0	13.4
80-84	10.1	8.7	22.1	6.7	5.8	15.3
85+	9.8	8.1	26.4	7.2	6.2	18.1

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.