

Table 10.15
Kaposi SarcomaEstimated United States Cancer Prevalence Counts^a on January 1, 2010
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <18 ^e	0 to <35 ^e	>=35 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	7,319	5,335	6,995	3,499	1,165	334	23,149	24,815	+	+
	Males	6,865	5,044	6,752	3,293	1,143	311	21,954	23,533	+	+
	Females	454	291	243	206	22	23	1,195	1,282	+	+
White ^b	Both Sexes	5,002	3,946	5,497	2,916	963	258	17,363	18,729	+	+
	Males	4,685	3,683	5,346	2,755	941	247	16,470	17,761	+	+
	Females	317	263	151	161	22	11	893	968	+	+
Black ^b	Both Sexes	1,915	1,174	1,068	374	93	40	4,531	4,676	+	+
	Males	1,814	1,151	1,022	352	93	40	4,339	4,484	+	+
	Females	101	23	46	22	0	0	192	192	+	+
Asian/ Pacific Islander ^c	Both Sexes	101	83	99	+	+	+	302	+	+	+
	Males	101	77	99	+	+	+	296	+	+	+
	Females	0	6	0	+	+	+	6	+	+	+
Hispanic ^d	Both Sexes	1,279	974	1,300	+	+	+	3,553	+	+	+
	Males	1,166	929	1,259	+	+	+	3,354	+	+	+
	Females	113	45	41	+	+	+	199	+	+	+

Estimated prevalence percent^a on January 1, 2010, of the SEER 11 population diagnosed in the previous 18 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	Age-Adjusted ^f All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0071%	-	-	0.0010%	0.0062%	0.0191%	0.0138%	0.0081%	0.0080%	0.0127%	0.0070%
	Males	0.0138%	-	-	0.0019%	0.0122%	0.0377%	0.0281%	0.0164%	0.0152%	0.0249%	0.0139%
	Females	0.0006%	-	-	-	0.0002%	0.0007%	0.0003%	0.0007%	0.0022%	0.0057%	0.0006%
White ^c	Both Sexes	0.0072%	-	-	0.0007%	0.0055%	0.0187%	0.0144%	0.0085%	0.0086%	0.0133%	0.0070%
	Males	0.0138%	-	-	0.0014%	0.0105%	0.0361%	0.0288%	0.0169%	0.0163%	0.0260%	0.0136%
	Females	0.0006%	-	-	-	-	0.0007%	-	0.0007%	0.0024%	0.0060%	0.0006%
Black ^c	Both Sexes	0.0103%	-	-	0.0032%	0.0148%	0.0322%	0.0173%	0.0102%	0.0042%	0.0098%	0.0109%
	Males	0.0209%	-	-	0.0065%	0.0313%	0.0674%	0.0365%	0.0211%	0.0088%	-	0.0227%
	Females	0.0007%	-	-	-	-	-	-	-	-	0.0082%	0.0008%
Asian/ Pacific Islander ^c	Both Sexes	0.0018%	-	-	-	0.0016%	0.0043%	0.0035%	0.0015%	0.0035%	0.0045%	0.0018%
	Males	0.0037%	-	-	-	0.0034%	0.0089%	0.0076%	0.0034%	0.0080%	0.0097%	0.0038%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	Both Sexes	0.0073%	-	-	0.0012%	0.0073%	0.0229%	0.0178%	0.0110%	0.0168%	0.0327%	0.0098%
	Males	0.0135%	-	-	0.0022%	0.0141%	0.0438%	0.0361%	0.0230%	0.0288%	0.0570%	0.0187%
	Females	0.0008%	-	-	-	-	0.0013%	-	-	0.0080%	0.0183%	0.0015%

^a US 2010 cancer prevalence counts are based on 2010 cancer prevalence proportions from the SEER registries and 1/1/2010 US population estimates based on the average of 2009 and 2010 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 35 years for 1975-2010 SEER 9 data; 18 years for 1992-2010 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 35 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <35 and >=35. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.