

Table 13.9
Chronic Myeloid Leukemia

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975-2010 | 1.76 | 2.33 | 1.34 | 1.78 | 2.37 | 1.36 | 1.77 | 2.33 | 1.37 |
| 1975 | 1.98 | 2.93 | 1.33 | 2.00 | 3.01 | 1.31 | 1.94 | - | - |
| 1976 | 1.93 | 2.48 | 1.55 | 1.90 | 2.45 | 1.52 | 1.86 | 2.58 | - |
| 1977 | 1.68 | 2.08 | 1.39 | 1.69 | 2.14 | 1.37 | 1.84 | - | - |
| 1978 | 1.71 | 2.26 | 1.34 | 1.74 | 2.28 | 1.39 | 1.43 | - | - |
| 1979 | 1.87 | 2.48 | 1.43 | 1.89 | 2.50 | 1.45 | 1.94 | 2.80 | - |
| 1980 | 1.70 | 2.06 | 1.46 | 1.73 | 2.10 | 1.51 | 1.48 | - | - |
| 1981 | 1.64 | 1.97 | 1.41 | 1.59 | 1.90 | 1.39 | 2.38 | 3.00 | 1.81 |
| 1982 | 1.68 | 2.16 | 1.35 | 1.65 | 2.19 | 1.31 | 1.75 | - | 1.79 |
| 1983 | 1.69 | 2.23 | 1.33 | 1.64 | 2.15 | 1.32 | 2.51 | 3.44 | 1.85 |
| 1984 | 1.67 | 2.40 | 1.18 | 1.62 | 2.40 | 1.10 | 2.58 | 2.61 | 2.59 |
| 1985 | 1.98 | 2.76 | 1.44 | 2.01 | 2.80 | 1.48 | 2.06 | 2.89 | - |
| 1986 | 1.68 | 2.20 | 1.29 | 1.71 | 2.26 | 1.31 | 1.54 | - | - |
| 1987 | 1.88 | 2.32 | 1.58 | 1.90 | 2.38 | 1.58 | 2.14 | 2.88 | - |
| 1988 | 1.91 | 2.54 | 1.50 | 1.93 | 2.60 | 1.49 | 2.08 | 2.88 | - |
| 1989 | 1.70 | 2.22 | 1.35 | 1.68 | 2.18 | 1.35 | 2.21 | 3.12 | - |
| 1990 | 1.76 | 2.47 | 1.24 | 1.75 | 2.50 | 1.20 | 2.63 | 3.66 | 1.95 |
| 1991 | 1.88 | 2.44 | 1.47 | 1.91 | 2.55 | 1.46 | 1.56 | - | 1.75 |
| 1992 | 1.76 | 2.16 | 1.47 | 1.80 | 2.22 | 1.49 | 1.49 | 1.73 | - |
| 1993 | 1.92 | 2.53 | 1.48 | 1.95 | 2.52 | 1.56 | 2.26 | 3.56 | - |
| 1994 | 1.86 | 2.47 | 1.42 | 1.90 | 2.48 | 1.50 | 1.70 | 2.57 | - |
| 1995 | 1.89 | 2.55 | 1.37 | 1.96 | 2.69 | 1.39 | 1.54 | 1.98 | - |
| 1996 | 1.81 | 2.45 | 1.35 | 1.81 | 2.53 | 1.27 | 1.99 | 2.39 | 1.83 |
| 1997 | 1.87 | 2.68 | 1.29 | 1.93 | 2.74 | 1.36 | 1.67 | 2.53 | - |
| 1998 | 1.65 | 1.95 | 1.42 | 1.73 | 1.97 | 1.55 | 1.48 | 2.31 | - |
| 1999 | 1.80 | 2.49 | 1.28 | 1.78 | 2.44 | 1.28 | 1.72 | 2.30 | 1.26 |
| 2000 | 1.78 | 2.33 | 1.36 | 1.80 | 2.33 | 1.40 | 1.55 | 1.72 | 1.33 |
| 2001 | 1.79 | 2.34 | 1.38 | 1.84 | 2.37 | 1.45 | 1.91 | 2.68 | 1.36 |
| 2002 | 1.61 | 2.05 | 1.28 | 1.63 | 2.08 | 1.31 | 1.10 | - | - |
| 2003 | 1.62 | 2.19 | 1.17 | 1.71 | 2.31 | 1.23 | 1.41 | 2.04 | - |
| 2004 | 1.78 | 2.51 | 1.22 | 1.85 | 2.62 | 1.25 | 1.47 | 1.97 | 1.11 |
| 2005 | 1.65 | 2.21 | 1.20 | 1.72 | 2.30 | 1.27 | 1.60 | 2.22 | 1.20 |
| 2006 | 1.80 | 2.38 | 1.36 | 1.90 | 2.50 | 1.45 | 1.61 | 2.17 | 1.16 |
| 2007 | 1.65 | 2.25 | 1.16 | 1.75 | 2.43 | 1.19 | 1.44 | 1.99 | 1.11 |
| 2008 | 1.74 | 2.23 | 1.34 | 1.73 | 2.16 | 1.40 | 1.73 | 2.42 | 1.11 |
| 2009 | 1.71 | 2.30 | 1.25 | 1.71 | 2.24 | 1.29 | 1.66 | 2.60 | 1.05 |
| 2010 | 1.70 | 2.29 | 1.23 | 1.64 | 2.26 | 1.15 | 1.76 | 2.01 | 1.64 |

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.