

Table 15.5  
Cancer of the Lung and Bronchus (Invasive)

Delay-adjusted SEER Incidence<sup>a</sup> Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975	52.26	89.56	24.53	51.96	89.16	24.85	64.55	114.90	24.66
1976	55.42	93.80	27.31	54.64	92.75	27.28	72.34	131.00	28.63
1977	56.70	95.51	28.31	55.88	94.76	28.04	73.58	124.86	33.88
1978	57.85	96.67	29.65	57.26	95.87	29.88	74.53	130.51	31.69
1979	58.62	95.81	31.59	58.04	95.19	31.65	74.54	127.72	34.11
1980	60.66	99.97	32.22	59.43	97.72	32.37	86.57	151.30	38.25
1981	62.03	99.41	35.12	61.54	98.44	35.52	82.31	142.96	37.45
1982	63.29	100.70	36.73	63.06	99.69	37.68	80.37	141.06	35.68
1983	63.45	99.46	37.89	63.04	97.63	39.10	86.62	151.34	39.49
1984	65.50	102.06	39.47	64.36	99.93	39.64	92.93	158.84	44.82
1985	64.63	98.64	40.24	64.00	96.90	40.89	89.56	149.69	46.05
1986	65.78	99.06	42.29	65.16	97.36	42.94	92.94	156.28	48.67
1987	67.90	101.52	44.16	68.08	100.65	45.50	84.81	143.72	43.28
1988	68.03	98.69	46.30	68.25	97.92	47.72	88.14	144.94	48.08
1989	67.54	97.95	46.09	67.56	97.05	47.24	89.05	142.78	51.37
1990	68.08	96.92	47.79	68.24	96.14	48.96	86.79	137.18	52.14
1991	69.21	97.23	49.59	68.99	95.92	50.64	93.16	144.78	57.06
1992	69.46	97.19	49.86	68.92	94.87	51.07	93.90	148.76	55.89
1993	67.77	93.97	49.18	67.84	92.53	50.67	86.33	135.38	52.45
1994	67.18	90.96	50.53	67.29	90.15	51.62	87.23	131.61	56.48
1995	66.86	89.83	50.45	67.12	87.99	52.46	85.73	138.76	50.02
1996	66.47	88.01	51.23	66.84	86.75	53.00	84.39	128.47	55.37
1997	66.64	86.30	52.62	67.27	84.88	55.00	82.17	129.25	50.52
1998	67.56	88.05	52.99	68.10	86.68	55.13	87.10	130.47	57.87
1999	65.87	84.62	52.53	66.37	83.36	54.49	84.46	122.01	59.00
2000	64.18	82.16	51.26	64.74	80.89	53.29	80.12	114.37	57.30
2001	64.21	81.44	51.73	65.12	81.11	53.59	80.35	114.07	57.61
2002	64.11	80.49	52.55	64.87	79.88	54.36	81.33	112.90	60.58
2003	64.82	81.10	53.15	65.66	79.74	55.77	80.70	118.62	55.49
2004	62.33	76.38	52.16	62.97	75.49	53.95	76.13	105.03	57.17
2005	63.14	75.97	53.92	64.21	75.79	55.95	74.66	97.88	59.52
2006	62.43	74.41	53.60	63.05	73.26	55.62	76.58	100.91	59.83
2007	62.17	73.78	53.70	63.34	73.03	56.31	74.14	100.23	56.87
2008	60.55	72.38	51.97	61.43	71.12	54.50	75.24	103.52	56.48
2009	60.03	70.78	52.27	60.83	69.98	54.22	73.72	97.20	58.25
2010	57.93	68.12	50.35	59.24	68.11	52.59	67.22	86.16	54.75

Additional information on the model used to delay-adjust SEER Incidence rates can be found at <http://surveillance.cancer.gov/delay/>.

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.