

Table 15.23
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted SEER Death Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Death Rates^a, 2006-2010</u>									
California	38.69	47.22	32.32	40.05	47.64	34.35	50.89	67.19	39.35
Greater Bay Area	35.12	41.63	30.40	36.11	40.55	32.91	55.31	70.81	44.45
San Francisco-Oakland	36.46	43.37	31.48	37.01	41.24	33.98	57.10	74.03	45.50
San Jose-Monterey	32.48	38.25	28.25	34.55	39.36	31.04	42.52	49.00	36.97
Los Angeles	34.51	44.46	27.20	34.18	42.36	28.09	51.20	69.80	39.10
Greater California	41.85	50.39	35.40	43.41	51.63	37.20	47.83	61.94	36.65
Connecticut	44.17	52.72	38.23	44.78	52.71	39.32	41.96	58.59	31.02
Detroit	55.37	69.81	45.32	54.51	67.21	45.73	61.79	84.61	46.83
Georgia	52.92	73.12	38.72	55.15	73.70	41.66	48.10	74.98	31.80
Atlanta	42.87	55.71	34.17	42.82	52.79	35.80	45.95	68.11	33.35
Rural Georgia	54.00	84.90	31.61	56.80	84.03	35.87	48.36	87.16	24.33
Greater Georgia	57.15	79.81	41.01	59.33	80.50	43.85	49.90	79.15	31.44
Hawaii	35.96	48.76	25.90	42.79	50.04	36.45	39.84	52.35	-
Iowa	49.33	64.06	38.34	49.29	63.94	38.37	71.93	97.23	52.04
Kentucky	73.17	97.05	55.80	73.19	96.70	55.97	80.75	114.09	59.08
Louisiana	59.40	80.90	43.63	58.45	76.12	45.32	63.66	98.45	40.32
New Jersey	44.79	55.78	37.25	46.27	56.72	39.09	46.30	65.48	34.47
New Mexico	35.11	43.36	28.57	36.62	44.78	30.14	48.24	73.98	25.54
Seattle-Puget Sound	47.41	55.62	41.37	48.62	56.19	43.03	54.12	65.29	44.77
Utah	21.60	27.54	16.76	21.43	27.27	16.67	53.38	68.38	-
9 SEER Areas ^b	42.89	52.99	35.49	43.56	52.49	37.00	53.44	73.54	40.20
11 SEER Areas ^b	40.45	50.29	33.23	41.17	49.75	34.84	52.74	72.23	39.87
13 SEER Areas ^b	40.64	50.54	33.36	41.33	49.95	34.95	52.63	72.36	39.63
18 SEER Areas ^b	45.40	57.27	36.64	46.57	57.45	38.49	53.41	76.45	38.15
Total U.S.	49.52	63.48	39.18	50.18	63.25	40.41	53.51	78.48	37.21

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta. The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.