


Table 15.25  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	63.48	0.10		
High Five States				
Kentucky	97.05	1.01	(01)	52.88 <sup>c</sup>
Mississippi	95.38	1.25	(02)	50.24 <sup>c</sup>
Arkansas	90.13	1.14	(03)	41.98 <sup>c</sup>
Tennessee	89.45	0.81	(04)	40.90 <sup>c</sup>
Alabama	87.37	0.91	(05)	37.63 <sup>c</sup>
Low Five States				
Hawaii	48.76	1.21	(47)	-23.20 <sup>c</sup>
California	47.22	0.26	(48)	-25.62 <sup>c</sup>
Colorado	44.20	0.70	(49)	-30.37 <sup>c</sup>
New Mexico	43.36	0.98	(50)	-31.70 <sup>c</sup>
Utah	27.54	0.80	(51)	-56.62 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	87.37	0.91	(05)	37.63 <sup>c</sup>	Montana	52.07	1.43	(43)	-17.98 <sup>c</sup>
Alaska	61.78	2.70	(27)	-2.68	Nebraska	60.33	1.20	(32)	-4.97
Arizona	49.82	0.58	(45)	-21.52 <sup>c</sup>	Nevada	59.66	1.06	(33)	-6.02 <sup>c</sup>
Arkansas	90.13	1.14	(03)	41.98 <sup>c</sup>	New Hampshire	59.00	1.40	(34)	-7.06 <sup>c</sup>
California	47.22	0.26	(48)	-25.62 <sup>c</sup>	New Jersey	55.78	0.53	(38)	-12.13 <sup>c</sup>
Colorado	44.20	0.70	(49)	-30.37 <sup>c</sup>	New Mexico	43.36	0.98	(50)	-31.70 <sup>c</sup>
Connecticut	52.72	0.79	(42)	-16.96 <sup>c</sup>	New York	54.31	0.35	(39)	-14.46 <sup>c</sup>
Delaware	68.19	1.80	(17)	7.42 <sup>c</sup>	North Carolina	76.61	0.63	(11)	20.68 <sup>c</sup>
D.C.	61.89	2.37	(26)	-2.51	North Dakota	54.14	1.80	(40)	-14.72 <sup>c</sup>
Florida	61.38	0.34	(29)	-3.32	Ohio	74.81	0.52	(13)	17.84 <sup>c</sup>
Georgia	73.12	0.68	(14)	15.18 <sup>c</sup>	Oklahoma	79.55	0.97	(08)	25.31 <sup>c</sup>
Hawaii	48.76	1.21	(47)	-23.20 <sup>c</sup>	Oregon	58.44	0.81	(35)	-7.94 <sup>c</sup>
Idaho	49.16	1.22	(46)	-22.56 <sup>c</sup>	Pennsylvania	65.83	0.45	(21)	3.70
Illinois	65.83	0.50	(20)	3.70	Rhode Island	64.92	1.61	(22)	2.26
Indiana	79.39	0.75	(09)	25.06 <sup>c</sup>	South Carolina	77.08	0.88	(10)	21.41 <sup>c</sup>
Iowa	64.06	0.91	(23)	0.90	South Dakota	61.91	1.76	(25)	-2.48
Kansas	66.77	1.02	(19)	5.18 <sup>c</sup>	Tennessee	89.45	0.81	(04)	40.90 <sup>c</sup>
Kentucky	97.05	1.01	(01)	52.88 <sup>c</sup>	Texas	60.71	0.38	(30)	-4.36
Louisiana	80.90	0.94	(07)	27.43 <sup>c</sup>	Utah	27.54	0.80	(51)	-56.62 <sup>c</sup>
Maine	69.94	1.41	(15)	10.16 <sup>c</sup>	Vermont	62.33	2.02	(24)	-1.81
Maryland	61.74	0.73	(28)	-2.75	Virginia	67.06	0.65	(18)	5.63 <sup>c</sup>
Massachusetts	60.61	0.64	(31)	-4.52	Washington	56.95	0.64	(37)	-10.29 <sup>c</sup>
Michigan	68.21	0.55	(16)	7.44 <sup>c</sup>	West Virginia	84.35	1.30	(06)	32.87 <sup>c</sup>
Minnesota	53.28	0.68	(41)	-16.07 <sup>c</sup>	Wisconsin	58.39	0.66	(36)	-8.03 <sup>c</sup>
Mississippi	95.38	1.25	(02)	50.24 <sup>c</sup>	Wyoming	49.90	2.03	(44)	-21.40 <sup>c</sup>
Missouri	76.49	0.74	(12)	20.48 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.