


Table 21.14
Cancer of the Ovary (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2006-2010

Females

| State | Rate | SE | Rank | PD |
|------------------|------|------|------|---------------------|
| TOTAL U.S. | 8.08 | 0.03 | | |
| High Five States | | | | |
| Washington | 9.56 | 0.23 | (01) | 18.23 ^c |
| Alabama | 9.24 | 0.25 | (02) | 14.37 |
| Oregon | 9.18 | 0.29 | (03) | 13.56 |
| Wisconsin | 9.08 | 0.23 | (04) | 12.34 |
| Iowa | 8.92 | 0.31 | (05) | 10.39 |
| Low Five States | | | | |
| Rhode Island | 7.30 | 0.46 | (47) | -9.67 |
| Mississippi | 7.27 | 0.29 | (48) | -10.01 |
| Nevada | 6.91 | 0.33 | (49) | -14.49 |
| Alaska | 6.66 | 0.76 | (50) | -17.59 ^c |
| Hawaii | 6.10 | 0.39 | (51) | -24.54 ^c |

| State | Rate | SE | Rank | PD | State | Rate | SE | Rank | PD |
|---------------|------|------|------|---------------------|----------------|------|------|------|--------------------|
| Alabama | 9.24 | 0.25 | (02) | 14.37 | Montana | 8.74 | 0.53 | (07) | 8.20 |
| Alaska | 6.66 | 0.76 | (50) | -17.59 ^c | Nebraska | 7.72 | 0.38 | (38) | -4.53 |
| Arizona | 7.77 | 0.21 | (35) | -3.87 | Nevada | 6.91 | 0.33 | (49) | -14.49 |
| Arkansas | 8.25 | 0.31 | (17) | 2.14 | New Hampshire | 7.95 | 0.45 | (32) | -1.58 |
| California | 8.03 | 0.09 | (27) | -0.67 | New Jersey | 8.71 | 0.18 | (08) | 7.72 |
| Colorado | 8.65 | 0.27 | (09) | 7.03 | New Mexico | 7.64 | 0.37 | (39) | -5.49 |
| Connecticut | 7.93 | 0.27 | (33) | -1.93 | New York | 8.06 | 0.12 | (26) | -0.24 |
| Delaware | 8.07 | 0.54 | (23) | -0.14 | North Carolina | 7.61 | 0.17 | (40) | -5.88 |
| D.C. | 7.49 | 0.69 | (44) | -7.33 | North Dakota | 8.37 | 0.65 | (15) | 3.52 |
| Florida | 7.34 | 0.11 | (46) | -9.13 | Ohio | 8.15 | 0.15 | (19) | 0.80 |
| Georgia | 8.37 | 0.19 | (14) | 3.63 | Oklahoma | 8.02 | 0.27 | (28) | -0.80 |
| Hawaii | 6.10 | 0.39 | (51) | -24.54 ^c | Oregon | 9.18 | 0.29 | (03) | 13.56 |
| Idaho | 7.78 | 0.44 | (34) | -3.78 | Pennsylvania | 8.79 | 0.14 | (06) | 8.72 |
| Illinois | 8.08 | 0.15 | (22) | -0.05 | Rhode Island | 7.30 | 0.46 | (47) | -9.67 |
| Indiana | 8.43 | 0.21 | (12) | 4.32 | South Carolina | 8.11 | 0.25 | (21) | 0.37 |
| Iowa | 8.92 | 0.31 | (05) | 10.39 | South Dakota | 7.96 | 0.57 | (30) | -1.46 |
| Kansas | 8.44 | 0.32 | (10) | 4.40 | Tennessee | 7.74 | 0.20 | (36) | -4.24 |
| Kentucky | 8.12 | 0.25 | (20) | 0.43 | Texas | 7.53 | 0.11 | (42) | -6.83 |
| Louisiana | 7.73 | 0.25 | (37) | -4.29 | Utah | 7.60 | 0.38 | (41) | -5.94 |
| Maine | 7.96 | 0.42 | (31) | -1.46 | Vermont | 7.52 | 0.63 | (43) | -6.98 |
| Maryland | 8.06 | 0.22 | (25) | -0.22 | Virginia | 8.43 | 0.20 | (11) | 4.33 |
| Massachusetts | 8.07 | 0.20 | (24) | -0.16 | Washington | 9.56 | 0.23 | (01) | 18.23 ^c |
| Michigan | 8.42 | 0.17 | (13) | 4.15 | West Virginia | 8.33 | 0.36 | (16) | 3.02 |
| Minnesota | 8.01 | 0.23 | (29) | -0.94 | Wisconsin | 9.08 | 0.23 | (04) | 12.34 |
| Mississippi | 7.27 | 0.29 | (48) | -10.01 | Wyoming | 8.17 | 0.74 | (18) | 1.13 |
| Missouri | 7.46 | 0.20 | (45) | -7.71 | | | | | |



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.