


Table 27.19
Cancer of the Urinary Bladder (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2006-2010

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.69	0.04		
High Five States				
Maine	10.54	0.56	(01)	37.02 ^c
Vermont	9.73	0.83	(02)	26.48 ^c
Massachusetts	9.22	0.25	(03)	19.90 ^c
Rhode Island	9.07	0.61	(04)	17.98 ^c
Ohio	9.04	0.19	(05)	17.56 ^c
Low Five States				
Mississippi	6.50	0.35	(47)	-15.45 ^c
Utah	6.40	0.40	(48)	-16.80 ^c
New Mexico	5.99	0.38	(49)	-22.17 ^c
Alaska	5.57	0.85	(50)	-27.61 ^c
Hawaii	4.78	0.38	(51)	-37.88 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.66	0.27	(44)	-13.40	Montana	8.28	0.59	(16)	7.61
Alaska	5.57	0.85	(50)	-27.61 ^c	Nebraska	6.95	0.41	(39)	-9.61
Arizona	6.95	0.23	(38)	-9.59	Nevada	8.77	0.44	(06)	14.09
Arkansas	7.34	0.34	(34)	-4.51	New Hampshire	8.73	0.56	(07)	13.58
California	6.83	0.10	(41)	-11.14	New Jersey	8.62	0.22	(08)	12.11
Colorado	6.73	0.28	(43)	-12.52	New Mexico	5.99	0.38	(49)	-22.17 ^c
Connecticut	8.31	0.32	(14)	8.11	New York	7.96	0.14	(21)	3.44
Delaware	8.30	0.65	(15)	7.94	North Carolina	7.36	0.21	(33)	-4.28
D.C.	6.52	0.79	(46)	-15.21 ^c	North Dakota	8.22	0.69	(17)	6.94
Florida	7.65	0.12	(26)	-0.57	Ohio	9.04	0.19	(05)	17.56 ^c
Georgia	7.51	0.24	(28)	-2.34	Oklahoma	7.55	0.31	(27)	-1.81
Hawaii	4.78	0.38	(51)	-37.88 ^c	Oregon	7.93	0.30	(22)	3.15
Idaho	7.43	0.49	(31)	-3.39	Pennsylvania	8.52	0.16	(10)	10.79
Illinois	7.87	0.18	(23)	2.37	Rhode Island	9.07	0.61	(04)	17.98 ^c
Indiana	8.07	0.25	(19)	4.90	South Carolina	6.94	0.28	(40)	-9.71
Iowa	8.34	0.33	(13)	8.40	South Dakota	7.03	0.60	(37)	-8.57
Kansas	6.63	0.33	(45)	-13.75	Tennessee	7.72	0.26	(25)	0.39
Kentucky	8.51	0.32	(11)	10.65	Texas	6.75	0.13	(42)	-12.19
Louisiana	7.23	0.30	(36)	-6.04	Utah	6.40	0.40	(48)	-16.80 ^c
Maine	10.54	0.56	(01)	37.02 ^c	Vermont	9.73	0.83	(02)	26.48 ^c
Maryland	8.34	0.28	(12)	8.43	Virginia	7.39	0.23	(32)	-3.88
Massachusetts	9.22	0.25	(03)	19.90 ^c	Washington	8.05	0.25	(20)	4.67
Michigan	8.58	0.20	(09)	11.52	West Virginia	7.76	0.41	(24)	0.90
Minnesota	7.24	0.26	(35)	-5.90	Wisconsin	8.21	0.26	(18)	6.80
Mississippi	6.50	0.35	(47)	-15.45 ^c	Wyoming	7.44	0.80	(30)	-3.27
Missouri	7.47	0.24	(29)	-2.87					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.