5-Year Relative Survival (Percent), 2003-2009^a

By International Classification of Childhood Cancer(ICCC)^b Selected Group and Subgroup^c and Sex and Age Excluding benign brain and myelodysplastic syndromes

Ages 0-19 by Sex		Botl	Both Sexes	Both Sexes by Age	Both Sexes by Age
Group & Subgroup Total Male Female	<1			1 3	
ICCC Groups Combined 82.1 81.1 83.2	77.5	77.5 83.8	77.5 83.8 81.7	77.5 83.8 81.7 80.9	77.5 83.8 81.7 80.9 82.8
eukemia81.480.782.4(a) Lymphoid leukemia86.685.288.6(b) Acute myeloid leukemia61.561.661.3	61.0 55.8 62.5	55.8 93.1	55.8 93.1 90.3	55.8 93.1 90.3 80.6	55.8 93.1 90.3 80.6 72.5
Lymphomas and reticuloendothelial neo. 91.2 91.0 91.6 I(a) Hodgkin lymphoma 96.6 97.2 95.9 I(b,c,e) Non-Hodgkin lymphoma 85.3 85.1 85.7	81.1 ^d _	- 96.5	- 96.5 96.4	- 96.5 96.4 97.1	- 96.5 96.4 97.1 96.4
CNS & misc intracranial & intraspinal neo.73.072.973.1II(a) Ependymomas and choroid plexus tumor75.571.180.7II(b) Astrocytoma84.083.984.1II(c) Intracranial and intraspinal embryonal tumors61.262.159.8II(d) Other gliomas57.461.153.7	56.6 59.0 ^d 82.2 25.7 ^d 75.3 ^d	59.0 ^d 74.5 82.2 91.0 25.7 ^d 55.5	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Neuroblastoma and other peripheral nervous cell tumors 78.5 77.7 79.4 V(a) Neuroblastoma and ganglioneuroblastoma 78.3 77.6 79.0	91.1 91.4				
etinoblastoma 97.5 96.1 99.0	99.9	99.9 95.6	99.9 95.6 -	99.9 95.6	99.9 95.6
Renal tumors 88.8 89.7 88.0 I(a) Nephroblastoma and other nonepithelial renal tumor 89.7 90.6 88.8	90.4 90.4				
Hepatic tumors 68.9 68.4 70.3 II(a) Hepatoblastoma 78.1 74.6 84.9	80.0 80.4				
I Malignant bone tumors 68.7 66.7 71.1 III(a) Osteosarcoma 66.6 64.2 69.4 III(c) Ewing tumor and related sarcomas of bone 67.3 64.2 71.3	- - -	,	66.7 ^d	66.7 ^d 72.4	66.7 ^d 72.4 60.7
Soft tissue and other extraosseous sarcomas70.569.371.8X(a) Rhabdomyosarcoma63.863.364.4	65.4 63.6 ^d				
erm cell & trophoblastic tumors & neo. of gonads 92.1 92.2 91.9 (a) Intracranial and intraspinal germ-cell tumor 90.1 91.8 84.0 ^d (c) Malignant gonadal germ cell tumor 95.4 94.6 97.1	92.1 		83.9 ^d	83.9 ^d 94.8	83.9 ^d 94.8 89.4
Other malignant epithelial neo. and melanomas ^f 92.8 88.2 95.1 I(b) Thyroid carcinoma 99.2 96.4 99.7 I(d) Malignant melanoma 95.7 93.0 97.5	- - -		98.1	98.1 100.0	98.1 100.0 99.0

^a SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b Based on follow-up of patients into 2010.
^b International Classification of Childhood Cancer is based on ICD-0-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005: Vol 103, No. 7, pg 1457-1467.
^c Classifications are shown for invasive cases only. For ICCC groups and subgroups that include *in situ* behavior,

only invasive cases are shown except as noted.

^d The standard error is between 5 and 10 percentage points.

- ^e The standard error is greater than 10 percentage points.
- f Includes in situ urinary bladder tumors.
- Statistic could not be calculated due to fewer than 25 cases during the time period. Neoplasms are abbreviated as neo.