Table 1.24
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites[a] by Race/Ethnicity

Both Sexes

All Races

Site	Rate[b] 2007-2011	APC[c] 2002-2011
All Sites	460.4	-0.6*
Breast	67.1	-0.3
Prostate[f]	67.0	-1.9*
Lung and Bronchus	60.1	-1.7*
Colon and Rectum	43.7	-2.8*
Melanoma of the Skin	21.3	1.4*
Urinary Bladder	20.5	-0.8*
Non-Hodgkin Lymphoma	19.7	-0.1
Kidney and Renal Pelvis	15.5	1.9*
Corpus and Uterus, NOS[f]	13.1	1.0*
Leukemia	13.0	0.0
Thyroid	12.9	5.8*
Pancreas	12.3	0.7*
Oral Cavity and Pharynx	11.0	0.4*
Liver & IBD[g]	7.9	3.4*
Stomach	7.5	-1.3*

White

Site	Rate[b] 2007-2011	APC[c] 2002-2011
All Sites	468.9	-0.6*
Breast	68.3	-0.5
Prostate[f]	64.2	-2.2*
Lung and Bronchus	61.7	-1.6*
Colon and Rectum	42.9	-2.9*
Melanoma of the Skin	25.2	1.3*
Urinary Bladder	22.4	-0.7*
Non-Hodgkin Lymphoma	20.6	-0.2
Kidney and Renal Pelvis	15.9	1.9*
Leukemia	13.7	0.0
Thyroid	13.7	5.8*
Corpus and Uterus, NOS[f]	13.4	0.8*
Pancreas	12.2	0.9*
Oral Cavity and Pharynx	11.4	0.8*
Brain and ONS[g]	7.1	-0.5
Liver & IBD[g]	7.0	4.1*

Black

Site	Rate[b] 2007-2011	APC[c] 2002-2011
All Sites	480.8	-1.0*
Prostate[f]	94.8	-2.0*
Breast	70.0	0.3
Lung and Bronchus	68.0	-1.9*
Colon and Rectum	53.6	-2.7*
Kidney and Renal Pelvis	17.9	2.7*
Pancreas	15.6	-0.2
Non-Hodgkin Lymphoma	14.3	-0.3
Corpus and Uterus, NOS[f]	13.3	2.3*
Urinary Bladder	12.6	-0.5
Myeloma	12.2	0.4
Stomach	11.2	-2.1*
Leukemia	10.0	-1.0
Liver & IBD[g]	9.4	3.4*
Oral Cavity and Pharynx	9.4	-2.5*
Thyroid	7.6	5.9*

Asian/Pacific Islander

Site	Rate[b] 2007-2011	APC[c] 2002-2011
All Sites	306.7	-0.9*
Breast	51.6	0.6
Lung and Bronchus	37.0	-1.4*
Colon and Rectum	36.9	-2.2*
Prostate[f]	34.5	-3.6*
Liver & IBD[g]	14.1	-0.9
Non-Hodgkin Lymphoma	13.4	0.0
Thyroid	12.5	5.5*
Stomach	11.5	-3.7*
Corpus and Uterus, NOS[f]	10.5	2.5*
Pancreas	9.7	0.9
Urinary Bladder	8.8	-1.6*
Kidney and Renal Pelvis	8.1	2.3*
Leukemia	7.6	-0.2
Oral Cavity and Pharynx	7.5	-1.1
Ovary[fh]	5.1	-1.5*

American Indian/Alaska Native[d]

Site	Rate[b] 2007-2011	APC[c] 2002-2011
All Sites	319.3	-0.8
Breast	42.8	0.1
Lung and Bronchus	40.9	-2.7*
Colon and Rectum	40.0	-1.7*
Prostate[f]	31.6	-3.6*
Kidney and Renal Pelvis	19.4	1.6
Liver & IBD[g]	13.7	3.9
Non-Hodgkin Lymphoma	12.2	1.8
Pancreas	10.7	0.6
Corpus and Uterus, NOS[f]	10.6	2.2
Stomach	9.6	-4.0*
Urinary Bladder	8.1	0.7
Oral Cavity and Pharynx	8.1	3.0
Thyroid	7.7	4.2*
Leukemia	7.3	0.1
Ovary[fh]	5.4	-2.1

Hispanic[e]

Site	Rate[b] 2007-2011	APC[c] 2002-2011
All Sites	353.2	-0.9*
Prostate[f]	53.6	-3.0*
Breast	49.1	-0.2
Colon and Rectum	36.6	-1.9*
Lung and Bronchus	31.3	-2.1*
Non-Hodgkin Lymphoma	17.7	0.3
Kidney and Renal Pelvis	15.2	1.7*
Liver & IBD[g]	12.2	2.8*
Urinary Bladder	11.4	-1.3*
Pancreas	11.1	-0.1
Stomach	11.1	-2.0*
Thyroid	10.9	5.3*
Corpus and Uterus, NOS[f]	10.6	1.7*
Leukemia	10.3	0.1
Oral Cavity and Pharynx	6.7	1.2*
Ovary[fh]	5.9	-2.1*

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

[a] Top 15 cancer sites selected based on 2007-2011 age-adjusted rates for the race/ethnic group.
[b] Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
[c] The APC is the Annual Percent Change over the time interval.
Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
[d] Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
[e] Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
[f] The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
[g] IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
[h] Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
* The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.