

Table 2.4
All Cancer Sites (Invasive)

Delay-adjusted SEER Incidence^a Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975	400.41	466.79	365.87	402.11	468.53	369.59	426.52	525.51	356.63
1976	407.38	481.11	367.41	409.71	481.86	373.14	427.45	552.95	340.63
1977	407.74	486.51	363.70	409.33	487.18	367.89	442.60	577.58	352.02
1978	407.29	488.35	362.20	409.81	489.71	368.02	436.25	570.67	343.64
1979	412.52	496.45	365.55	414.16	497.62	369.74	451.53	589.70	359.09
1980	418.08	505.84	368.13	419.50	505.22	373.38	467.48	615.10	366.55
1981	425.32	510.91	377.17	427.73	510.89	383.94	472.73	633.91	362.94
1982	424.54	511.11	376.27	427.85	512.55	383.53	465.18	621.92	359.41
1983	431.20	520.01	380.81	434.48	521.08	388.07	487.09	646.07	382.06
1984	440.09	525.55	392.54	443.80	527.86	399.72	488.25	643.94	386.08
1985	449.01	531.38	403.31	453.67	533.80	411.66	489.19	643.65	388.43
1986	451.54	538.43	402.45	456.26	543.20	409.37	493.98	644.19	396.60
1987	468.20	563.42	413.42	475.14	569.18	423.49	501.24	666.32	394.36
1988	464.06	558.64	409.43	470.59	565.46	418.12	504.12	657.88	404.49
1989	468.04	567.33	408.57	475.53	574.00	418.89	493.00	661.75	384.86
1990	482.30	593.07	415.84	489.37	599.91	425.21	522.27	700.07	409.49
1991	503.38	637.86	419.20	511.05	645.21	428.96	549.39	765.32	412.79
1992	510.98	658.65	415.15	517.44	664.19	423.97	569.12	809.33	413.59
1993	493.82	622.65	408.64	497.60	621.06	417.61	568.79	819.04	404.34
1994	483.93	590.23	413.93	488.09	588.37	423.67	558.52	784.25	409.47
1995	477.44	572.28	415.24	482.92	571.08	426.78	543.52	750.28	405.00
1996	479.77	573.26	418.60	485.77	575.07	428.88	541.24	727.97	417.57
1997	487.01	576.31	428.91	494.52	577.65	441.95	541.16	741.14	410.61
1998	488.93	573.64	434.42	497.87	577.73	447.84	536.79	723.99	416.22
1999	491.26	581.57	431.20	499.84	584.25	444.70	541.35	724.63	421.10
2000	487.07	581.17	422.84	497.87	585.67	438.94	528.43	713.55	404.66
2001	490.83	582.36	427.85	502.52	589.34	443.65	521.44	699.37	401.59
2002	488.66	575.86	428.98	499.38	583.23	442.40	535.71	699.65	426.35
2003	476.44	561.26	418.20	487.85	567.20	433.95	520.13	681.39	412.84
2004	477.66	559.54	421.01	488.67	566.26	435.17	525.56	685.98	419.94
2005	473.81	550.57	420.58	486.68	560.43	435.90	509.88	644.31	421.02
2006	477.47	557.57	420.36	490.30	566.35	436.21	509.57	645.41	416.66
2007	483.56	568.67	422.11	495.56	575.99	437.67	519.10	668.88	416.31
2008	477.22	550.03	425.44	488.75	556.68	440.71	518.74	665.06	419.48
2009	475.89	540.78	429.95	487.35	548.17	444.41	511.46	640.47	424.14
2010	466.64	530.53	420.82	478.57	537.79	436.39	494.32	617.66	409.31
2011	463.44	520.77	423.17	476.22	529.27	439.39	490.50	596.02	418.33

Additional information on the model used to delay-adjust SEER Incidence rates can be found at [\(http://surveillance.cancer.gov/delay/\)](http://surveillance.cancer.gov/delay/).

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.