

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2007-2011</u>						
Age at Diagnosis/Death:						
All ages	7.8	7.8	9.4	2.3	2.1	4.1
Under 65	7.3	7.5	7.7	1.8	1.7	2.8
65 and over	11.6	10.3	20.8	6.0	5.2	12.9
All ages (IARC world std) ^c	6.2	6.2	7.1	1.7	1.6	2.9
<u>Age-Specific Rates, 2007-2011</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	0.1	-	-	-	-	-
20-24	1.3	1.4	1.3	0.1	0.1	0.2
25-29	5.1	5.6	4.2	0.5	0.5	0.8
30-34	10.1	11.1	9.1	1.3	1.2	1.8
35-39	13.9	15.3	10.5	2.3	2.3	3.3
40-44	14.8	15.3	15.2	3.2	3.0	4.6
45-49	13.8	13.8	14.7	4.0	3.7	6.0
50-54	12.9	12.4	16.5	4.5	4.1	7.3
55-59	12.1	11.4	17.3	4.7	4.3	7.4
60-64	12.6	11.5	17.8	5.2	4.8	9.2
65-69	12.8	12.0	20.1	5.5	4.9	11.3
70-74	12.5	11.1	18.9	5.6	5.0	11.2
75-79	10.8	9.7	19.9	5.8	5.0	13.2
80-84	10.1	8.8	21.5	6.7	5.9	15.1
85+	9.8	7.9	26.9	7.0	6.1	16.6

^a SEER 18 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.