

Table 6.24
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted SEER Death Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Death Rates^a, 2007-2011</u>									
California	14.53	17.19	12.43	14.53	17.24	12.37	21.99	26.10	19.05
Greater Bay Area	13.98	16.36	12.09	14.10	16.63	12.05	22.04	26.46	19.29
San Francisco-Oakland	14.63	17.40	12.47	14.71	17.75	12.24	22.19	26.92	19.19
San Jose-Monterey	12.65	14.34	11.29	12.98	14.64	11.63	20.92	24.36	19.21
Los Angeles	14.86	17.94	12.55	14.48	17.57	12.13	22.53	26.48	19.74
Greater California	14.56	17.13	12.49	14.67	17.29	12.55	21.27	25.40	18.04
Connecticut	13.11	15.28	11.42	13.01	15.12	11.37	15.68	18.91	13.23
Detroit	17.01	20.42	14.64	15.62	18.65	13.45	22.93	29.04	19.11
Georgia	16.12	19.83	13.49	14.57	18.06	11.97	21.77	27.56	18.47
Atlanta	15.31	18.71	13.04	13.05	15.98	10.96	20.79	26.59	17.65
Rural Georgia	17.50	24.70	12.91	14.56	20.57	10.25	22.91	33.19	17.22
Greater Georgia	16.45	20.18	13.73	15.14	18.73	12.41	22.35	27.89	19.07
Hawaii	13.78	17.59	10.66	13.37	15.71	10.94	-	-	-
Iowa	17.07	19.96	14.83	17.01	19.92	14.74	24.94	27.34	24.04
Kentucky	18.92	22.60	16.05	18.62	22.29	15.77	25.77	29.79	22.21
Louisiana	18.87	23.33	15.37	16.91	20.69	13.81	25.30	33.08	20.04
New Jersey	16.81	20.25	14.29	16.59	20.02	14.04	22.30	28.05	18.67
New Mexico	15.25	18.66	12.47	15.31	18.99	12.32	22.11	-	30.30
Seattle-Puget Sound	13.96	16.18	12.13	13.97	16.13	12.14	19.80	22.70	17.48
Utah	12.06	14.05	10.33	11.99	13.94	10.28	-	-	-
9 SEER Areas ^b	14.88	17.64	12.74	14.51	17.15	12.41	21.18	26.01	18.10
11 SEER Areas ^b	14.75	17.51	12.62	14.42	17.10	12.32	21.55	26.13	18.55
13 SEER Areas ^b	14.78	17.56	12.63	14.42	17.09	12.30	21.53	26.19	18.49
18 SEER Areas ^b	15.54	18.53	13.22	15.19	18.10	12.89	22.40	27.67	18.91
Total U.S.	15.94	19.12	13.46	15.47	18.55	13.02	22.14	27.67	18.51

^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta. The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

- Statistic not shown. Rate based on less than 16 cases for the time interval.