


Table 6.27  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2007-2011

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	13.46	0.04		
High Five States				
D.C.	18.77 <sup>b</sup>	1.07	(01)	39.47 <sup>c</sup>
Mississippi	16.50 <sup>b</sup>	0.44	(02)	22.59 <sup>c</sup>
Kentucky	16.05 <sup>b</sup>	0.35	(03)	19.23 <sup>c</sup>
West Virginia	15.62 <sup>b</sup>	0.49	(04)	16.01 <sup>c</sup>
Arkansas	15.61 <sup>b</sup>	0.42	(05)	15.95 <sup>c</sup>
Low Five States				
Colorado	12.03 <sup>b</sup>	0.31	(47)	-10.63 <sup>c</sup>
Arizona	11.68 <sup>b</sup>	0.25	(48)	-13.20 <sup>c</sup>
Connecticut	11.42 <sup>b</sup>	0.31	(49)	-15.18 <sup>c</sup>
Hawaii	10.66 <sup>b</sup>	0.50	(50)	-20.84 <sup>c</sup>
Utah	10.33 <sup>b</sup>	0.44	(51)	-23.25 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	14.42	0.31	(14)	7.14	Montana	12.72	0.64	(37)	-5.53
Alaska	13.88	1.13	(20)	3.10	Nebraska	14.97	0.51	(08)	11.22 <sup>c</sup>
Arizona	11.68 <sup>b</sup>	0.25	(48)	-13.20 <sup>c</sup>	Nevada	14.29	0.47	(15)	6.18
Arkansas	15.61 <sup>b</sup>	0.42	(05)	15.95 <sup>c</sup>	New Hampshire	13.25	0.57	(31)	-1.54
California	12.43 <sup>b</sup>	0.11	(43)	-7.64	New Jersey	14.29 <sup>b</sup>	0.23	(16)	6.14
Colorado	12.03 <sup>b</sup>	0.31	(47)	-10.63 <sup>c</sup>	New Mexico	12.47	0.46	(41)	-7.34
Connecticut	11.42 <sup>b</sup>	0.31	(49)	-15.18 <sup>c</sup>	New York	13.33	0.15	(29)	-0.97
Delaware	13.47	0.69	(27)	0.07	North Carolina	12.69 <sup>b</sup>	0.21	(39)	-5.70
D.C.	18.77 <sup>b</sup>	1.07	(01)	39.47 <sup>c</sup>	North Dakota	13.14	0.77	(33)	-2.39
Florida	12.45 <sup>b</sup>	0.14	(42)	-7.54	Ohio	14.61 <sup>b</sup>	0.20	(11)	8.51
Georgia	13.49	0.24	(26)	0.22	Oklahoma	14.28	0.36	(17)	6.06
Hawaii	10.66 <sup>b</sup>	0.50	(50)	-20.84 <sup>c</sup>	Oregon	13.30	0.34	(30)	-1.17
Idaho	12.30	0.55	(45)	-8.60	Pennsylvania	14.55 <sup>b</sup>	0.18	(12)	8.10
Illinois	14.83 <sup>b</sup>	0.20	(09)	10.20 <sup>c</sup>	Rhode Island	13.41	0.61	(28)	-0.40
Indiana	14.25	0.27	(18)	5.82	South Carolina	13.68	0.31	(23)	1.61
Iowa	14.83 <sup>b</sup>	0.38	(10)	10.17 <sup>c</sup>	South Dakota	13.61	0.71	(24)	1.07
Kansas	13.08	0.39	(34)	-2.80	Tennessee	15.25 <sup>b</sup>	0.28	(07)	13.29 <sup>c</sup>
Kentucky	16.05 <sup>b</sup>	0.35	(03)	19.23 <sup>c</sup>	Texas	12.83 <sup>b</sup>	0.15	(36)	-4.69
Louisiana	15.37 <sup>b</sup>	0.35	(06)	14.21 <sup>c</sup>	Utah	10.33 <sup>b</sup>	0.44	(51)	-23.25 <sup>c</sup>
Maine	13.73	0.54	(21)	1.99	Vermont	14.48	0.85	(13)	7.58
Maryland	13.19	0.28	(32)	-2.03	Virginia	13.52	0.25	(25)	0.44
Massachusetts	12.70	0.24	(38)	-5.68	Washington	12.57	0.26	(40)	-6.62
Michigan	13.73	0.21	(22)	1.98	West Virginia	15.62 <sup>b</sup>	0.49	(04)	16.01 <sup>c</sup>
Minnesota	12.12 <sup>b</sup>	0.28	(46)	-9.94	Wisconsin	12.42 <sup>b</sup>	0.26	(44)	-7.73
Mississippi	16.50 <sup>b</sup>	0.44	(02)	22.59 <sup>c</sup>	Wyoming	12.84	0.93	(35)	-4.60
Missouri	14.09	0.27	(19)	4.64					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.