

Table 8.8
Cancer of the Esophagus (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	4	4	6	1	0	2
1970-1973 ^a	-	-	-	4	4	4	4	4	3
1975-1977 ^b	5.0	4.0	7.5	5.5	4.8	7.0	3.5	2.0	7.7
1978-1980 ^b	5.0	4.8	5.6	5.3	5.3	5.2	4.3	3.4	6.8
1981-1983 ^b	6.6	6.0	8.2	7.3	6.5	9.0	4.3	3.7	6.5
1984-1986 ^b	9.5	8.4	12.1	10.2	9.0	13.0	8.7	8.2	9.9
1987-1989 ^b	9.4	9.4	9.5	10.5	11.0	9.2	6.6	5.3	10.2
1990-1992 ^b	12.1	11.7	13.4	12.9	12.3	14.8	9.1	9.4	8.4
1993-1995 ^b	12.1	12.2	11.7	13.4	13.6	12.7	7.6	7.8	6.9
1996-1998 ^b	13.4	13.0	14.4	14.2	14.0	14.7	10.2	8.5	13.6
1999-2003 ^b	17.7	17.4	18.8	18.7	18.7	18.7	12.0	9.5	17.3
2004-2010 ^b	20.0 ^f	19.9 ^f	20.0 ^f	21.0 ^f	20.9 ^f	21.5 ^f	13.2 ^f	12.8 ^f	13.9 ^f
5-Year Period Survival (Percent) ^{cd}									
2010	17.9	18.2	16.7	18.6	19.0	17.0	11.9	10.8	14.3
Stage Distribution (%) 2004-2010 ^{ce}									
All Stages									
Number of cases	20,656	16,155	4,501	17,121	13,640	3,481	2,344	1,627	717
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	21	21	24	22	21	24	19	18	22
Regional	30	30	29	30	30	28	32	32	32
Distant	37	39	31	37	39	31	37	39	32
Unstaged	11	10	17	11	10	17	12	11	13
5-Year Relative Survival (Percent), 2004-2010 ^c									
Age at Diagnosis:									
Ages <45	20.0	21.1	15.4	20.6	21.5	15.2	15.6	16.4 ^g	15.3 ^g
Ages 45-54	20.0	18.8	25.8	21.8	20.2	30.9	10.0	8.5	14.2
Ages 55-64	19.7	19.3	21.8	20.8	20.5	22.4	13.6	11.8	18.7
Ages 65-74	19.3	19.3	19.4	20.1	20.1	19.9	12.3	11.3	14.8
Ages 75+	11.6	11.9	10.9	12.0	12.1	11.2	7.5	8.4	6.0
Ages <65	19.8	19.3	22.6	21.1	20.5	24.6	12.5	11.0	16.7
Ages 65+	15.5	16.0	14.3	16.0	16.5	14.6	10.3	10.2	10.5
Stage ^e :									
All Stages	17.5	17.6	17.2	18.3	18.4	17.8	11.6	10.7	13.7
Localized	39.6	41.4	33.8	41.8	43.7	35.4	22.7	21.5	24.8
Regional	21.1	21.4	19.7	21.9	22.2	20.8	14.6	15.0	14.1
Distant	3.8	3.6	4.8	3.8	3.6	4.8	3.3	2.3	6.2
Unstaged	11.5	11.3	11.9	11.1	11.2	10.8	11.5	10.3	14.0 ^g

^a Based on End Results data from a series of hospital registries and one population-based registry.

^b SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Based on follow-up of patients into 2011.

^c SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Based on follow-up of patients into 2011.

^d Period survival provides a 2010 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2007-2009: 0-1 year survival), (2006-2008: 1-2 year survival), (2005-2007: 2-3 year survival), (2004-2006: 3-4 year survival), (2003-2005: 4-5 years survival).

^e Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.

^f The difference between 1975-1977 and 2004-2010 is statistically significant ($p < .05$).

^g The standard error is between 5 and 10 percentage points.

^h The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.