

Table 8.9
Cancer of the Esophagus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	26.0	29.9	34.9	38.5	40.3	42.0	40.1	40.8	44.3	46.1	46.3	41.9	46.6	48.2	45.6	47.6	50.0	47.7	48.7	49.5
2-year	11.4	14.3	19.0	21.8	22.6	24.3	23.9	23.9	26.9	28.4	29.4	25.9	30.4	30.1	31.2	30.5	32.9	32.2	30.5	
3-year	7.7	10.0	13.5	16.3	16.8	18.6	17.5	18.3	20.9	23.9	23.1	20.8	23.8	25.8	24.2	25.1	26.9	24.4		
4-year	5.9	8.2	10.8	13.7	13.2	15.9	14.3	14.2	18.4	20.6	19.2	17.8	20.0	22.2	21.4	21.8	23.3			
5-year	4.9	7.0	9.5	12.2	11.6	15.1	12.4	12.6	17.5	18.9	17.9	16.8	17.2	20.2	18.9	20.0				
6-year	4.3	6.1	8.5	11.1	10.8	13.9	11.3	11.6	16.0	18.0	15.9	15.2	14.4	19.7	17.4					
7-year	3.8	5.6	7.7	10.2	10.2	12.9	10.7	11.3	14.8	16.7	14.8	14.8	13.1	18.3						
8-year	3.4	4.9	7.0	9.5	9.6	12.2	9.8	9.8	14.3	15.8	13.9	13.6	12.8							
9-year	3.2	4.5	6.4	9.0	9.0	11.6	9.3	9.6	13.6	15.7	13.6	13.1								
10-year	2.9	4.3	6.0	8.3	8.6	10.7	9.2	9.5	13.0	15.1	13.3									
11-year	2.7	3.9	5.4	7.8	7.7	9.7	9.0	9.1	12.8	14.3										
12-year	2.6	3.8	4.9	7.1	7.2	9.4	7.5	8.6	11.9											
13-year	2.4	3.5	4.6	7.0	6.8	9.4	6.9	8.6												
14-year	2.4	3.4	4.3	6.6	5.7	9.0	6.3													
15-year	2.4	3.4	4.2	6.2	5.5	8.7														
16-year	2.3	3.2	3.7	5.9	4.9															
17-year	2.3	3.0	3.2	5.5																
18-year	2.2	2.7	3.1																	
19-year	2.2	2.5	2.9																	
20-year	2.1	2.4	2.6																	

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).