

Table 8.10
Cancer of the Esophagus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	23.9	29.2	34.1	38.9	39.9	43.2	40.6	41.0	43.6	47.5	47.2	43.2	46.2	47.5	46.7	48.5	51.7	48.7	49.9	48.3
2-year	10.0	13.8	18.4	21.6	22.8	24.8	24.7	24.3	26.4	28.7	29.7	26.5	29.9	30.0	31.1	30.9	33.2	32.8	30.7	
3-year	6.8	9.2	13.1	16.4	16.7	18.2	18.3	18.5	20.7	23.6	23.2	21.3	23.1	25.6	24.1	25.9	27.4	24.9		
4-year	5.1	7.3	10.3	13.4	13.0	15.1	15.2	14.2	18.2	19.6	19.4	18.1	19.5	22.1	20.9	22.9	23.5			
5-year	4.2	6.0	9.3	12.0	11.8	13.9	13.0	12.1	17.3	17.3	18.4	16.9	16.7	19.7	18.5	21.1				
6-year	3.7	5.1	8.4	10.8	11.0	12.9	11.8	11.5	16.2	16.5	16.3	15.5	13.8	19.1	17.2					
7-year	3.3	4.5	7.6	10.2	10.7	11.9	11.5	11.2	15.1	15.4	15.4	15.3	12.6	18.1						
8-year	3.0	4.0	6.9	9.3	9.9	11.3	10.3	9.4	14.8	14.6	14.7	13.7	12.1							
9-year	2.8	3.6	6.5	8.8	9.6	10.6	10.0	9.3	14.2	14.4	14.2	13.3								
10-year	2.7	3.4	6.2	7.9	9.3	10.0	9.7	9.3	13.4	14.1	13.9									
11-year	2.5	3.3	5.7	7.5	8.4	9.0	9.5	9.1	13.3	13.1										
12-year	2.4	3.2	5.1	6.8	7.7	8.7	8.4	8.8	12.8											
13-year	2.3	3.0	4.6	6.8	7.1	8.7	7.8	8.8												
14-year	2.2	3.0	4.2	6.6	6.3	8.3	6.9													
15-year	2.2	3.0	4.0	6.2	5.9	8.3														
16-year	2.2	2.8	3.4	6.0	5.3															
17-year	2.2	2.8	2.9	5.6																
18-year	2.2	2.5	2.8																	
19-year	2.1	2.3	2.5																	
20-year	2.0	2.2	2.3																	

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).