

Table 8.11
Cancer of the Esophagus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	31.2	31.6	37.1	37.3	41.4	38.5	38.8	39.9	46.6	41.6	42.9	37.9	48.1	50.4	41.7	44.7	44.3	44.0	44.4	54.2
2-year	15.1	15.6	20.7	22.4	22.2	23.0	21.6	22.5	28.6	27.4	27.9	24.0	32.4	30.5	31.5	29.0	31.9	29.7	30.0	
3-year	10.0	12.0	14.5	16.2	17.0	19.9	15.1	17.6	21.3	24.6	22.7	19.2	26.3	26.5	24.4	22.0	25.4	22.2		
4-year	7.9	10.5	12.3	14.5	13.7	18.2	11.7	14.3	18.9	23.8	18.6	17.0	21.8	22.5	23.4	17.9	22.3			
5-year	6.5	9.3	10.0	12.9	10.7	18.2	10.6	14.1	17.9	23.8	16.1	16.6	18.7	21.5	20.4	16.1				
6-year	5.9	8.6	8.5	11.8	9.9	16.4	9.8	12.0	15.2	22.7	14.2	14.4	16.3	21.2	18.1					
7-year	5.2	8.3	8.0	10.2	9.0	15.8	8.1	11.7	13.4	20.7	12.3	13.0	15.0	18.6						
8-year	4.4	7.1	7.3	10.0	8.7	14.8	8.1	10.9	12.1	19.7	10.7	12.7	15.0							
9-year	4.0	6.5	6.3	9.5	7.1	14.2	7.4	10.5	11.0	19.3	10.7	11.8								
10-year	3.5	6.5	5.3	9.2	6.6	12.8	7.4	9.6	11.0	18.3	10.7									
11-year	3.3	5.4	4.8	8.8	5.6	11.9	7.3	9.3	10.1	17.7										
12-year	3.1	5.2	4.6	7.9	5.6	11.4	4.8	7.6	8.2											
13-year	2.8	4.7	4.6	7.1	5.6	11.4	4.2	7.6												
14-year	2.8	4.5	4.5	6.4	3.9	10.8	4.2													
15-year	2.8	4.5	4.5	6.1	3.9	8.8														
16-year	2.7	3.8	4.2	5.5	3.4															
17-year	2.4	3.2	3.9	5.4																
18-year	2.2	3.0	3.8																	
19-year	2.2	2.8	3.7																	
20-year	2.2	2.8	3.3																	

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).