

Table 8.12
Cancer of the Esophagus (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.50	0.49	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.34	0.30
	10	0.00	0.00	0.00	0.51	0.50		10	0.00	0.00	0.00	0.35	0.30
	20	0.00	0.00	0.02	0.51	0.50		20	0.00	0.00	0.01	0.35	0.30
	30	0.00	0.02	0.08	0.52	0.50		30	0.00	0.01	0.04	0.35	0.30
	40	0.02	0.08	0.22	0.52	0.51		40	0.01	0.03	0.10	0.35	0.30
	50	0.07	0.21	0.37	0.52	0.50		50	0.03	0.09	0.19	0.35	0.30
	60	0.15	0.33	0.45	0.48	0.47		60	0.07	0.16	0.26	0.33	0.29
	70	0.20	0.34	-	0.37	0.38		70	0.10	0.20	-	0.28	0.26
White	80	0.18	-	-	0.23	0.26	American Indian/ Alaska Native ^a	80	0.12	-	-	0.21	0.21
	0	0.00	0.00	0.00	0.53	0.51		0	0.00	0.00	0.00	0.32	0.35
	10	0.00	0.00	0.00	0.53	0.52		10	0.00	0.00	0.01	0.32	0.36
	20	0.00	0.00	0.02	0.54	0.52		20	0.00	0.01	0.02	0.33	0.36
	30	0.00	0.02	0.09	0.54	0.52		30	0.01	0.02	0.09	0.33	0.37
	40	0.02	0.09	0.23	0.54	0.52		40	0.02	0.08	0.15	0.33	0.37
	50	0.07	0.22	0.39	0.54	0.52		50	0.07	0.14	0.23	0.33	0.38
	60	0.16	0.34	0.47	0.50	0.49		60	0.08	0.17	0.24	0.27	0.37
Black	70	0.21	0.35	-	0.38	0.40	Hispanic ^b	70	0.11	0.19	-	0.23	0.32
	80	0.19	-	-	0.23	0.27		80	0.12	-	-	0.17	0.27
	0	0.00	0.00	0.00	0.45	0.40		0	0.00	0.00	0.00	0.38	0.35
	10	0.00	0.00	0.00	0.46	0.41		10	0.00	0.00	0.00	0.38	0.35
	20	0.00	0.00	0.02	0.46	0.41		20	0.00	0.00	0.01	0.38	0.35
	30	0.00	0.02	0.10	0.46	0.42		30	0.00	0.01	0.05	0.38	0.35
	40	0.02	0.10	0.24	0.47	0.42		40	0.01	0.05	0.13	0.38	0.35
	50	0.09	0.23	0.38	0.47	0.43		50	0.04	0.12	0.24	0.38	0.35
	60	0.16	0.32	0.40	0.42	0.39		60	0.09	0.22	0.33	0.36	0.34
	70	0.19	0.30	-	0.32	0.30		70	0.14	0.27	-	0.30	0.30
	80	0.15	-	-	0.19	0.19		80	0.16	-	-	0.20	0.24

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.