

Table 8.13
Cancer of the Esophagus (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.80	0.79	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.53	0.47
	10	0.00	0.00	0.00	0.81	0.80		10	0.00	0.00	0.00	0.54	0.48
	20	0.00	0.00	0.03	0.82	0.80		20	0.00	0.00	0.01	0.54	0.48
	30	0.00	0.03	0.14	0.82	0.81		30	0.00	0.01	0.06	0.54	0.48
	40	0.03	0.14	0.37	0.83	0.82		40	0.01	0.06	0.18	0.54	0.48
	50	0.11	0.35	0.62	0.83	0.83		50	0.05	0.17	0.32	0.54	0.48
	60	0.26	0.55	0.73	0.77	0.78		60	0.12	0.28	0.41	0.51	0.45
	70	0.34	0.55	-	0.60	0.64		70	0.17	0.32	-	0.42	0.40
White	80	0.31	-	-	0.38	0.44	American Indian/ Alaska Native ^a	80	0.18	-	-	0.31	0.32
	0	0.00	0.00	0.00	0.85	0.83		0	0.00	0.00	0.00	0.49	0.48
	10	0.00	0.00	0.01	0.86	0.84		10	0.00	0.00	0.01	0.50	0.49
	20	0.00	0.01	0.04	0.86	0.84		20	0.00	0.01	0.03	0.51	0.50
	30	0.00	0.04	0.15	0.87	0.85		30	0.01	0.03	0.14	0.52	0.51
	40	0.03	0.14	0.39	0.88	0.86		40	0.02	0.14	0.23	0.53	0.52
	50	0.12	0.37	0.66	0.88	0.86		50	0.12	0.22	0.33	0.53	0.54
	60	0.27	0.58	0.77	0.82	0.81		60	0.11	0.23	0.37	0.44	0.52
Black	70	0.36	0.58	-	0.63	0.66	Hispanic ^b	70	0.14	0.30	-	0.39	0.43
	80	0.32	-	-	0.39	0.45		80	0.23	-	-	0.36	0.34
	0	0.00	0.00	0.00	0.63	0.62		0	0.00	0.00	0.00	0.61	0.58
	10	0.00	0.00	0.00	0.65	0.63		10	0.00	0.00	0.00	0.62	0.58
	20	0.00	0.00	0.03	0.65	0.64		20	0.00	0.00	0.02	0.62	0.59
	30	0.00	0.03	0.14	0.67	0.65		30	0.00	0.02	0.08	0.63	0.59
	40	0.03	0.14	0.35	0.68	0.66		40	0.02	0.08	0.22	0.63	0.60
	50	0.12	0.34	0.56	0.69	0.67		50	0.06	0.21	0.43	0.63	0.60
	60	0.25	0.49	0.62	0.63	0.63		60	0.15	0.39	0.56	0.60	0.58
	70	0.30	0.47	-	0.49	0.50		70	0.26	0.46	-	0.50	0.52
	80	0.27	-	-	0.30	0.33		80	0.26	-	-	0.32	0.41

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.