


Table 8.18  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
TOTAL U.S.	4.22	0.02							
High Five States									
New Hampshire	5.73 <sup>b</sup>	0.28	(01)	35.83 <sup>c</sup>					
Maine	5.69 <sup>b</sup>	0.26	(02)	34.94 <sup>c</sup>					
Rhode Island	5.12	0.29	(03)	21.56 <sup>c</sup>					
Ohio	5.02 <sup>b</sup>	0.09	(04)	19.15 <sup>c</sup>					
Indiana	4.99 <sup>b</sup>	0.12	(05)	18.28 <sup>c</sup>					
Low Five States									
Texas	3.49 <sup>b</sup>	0.06	(47)	-17.21 <sup>c</sup>					
California	3.46 <sup>b</sup>	0.04	(48)	-17.97 <sup>c</sup>					
Hawaii	3.23 <sup>b</sup>	0.20	(49)	-23.38 <sup>c</sup>					
New Mexico	3.19 <sup>b</sup>	0.17	(50)	-24.22 <sup>c</sup>					
Utah	2.87 <sup>b</sup>	0.17	(51)	-31.86 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.