

Table 10.15
Kaposi SarcomaEstimated United States Cancer Prevalence Counts^a on January 1, 2011
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <19 ^e	0 to <36 ^e	>=36 ^g	Complete ^h
<u>Race</u>	<u>Sex</u>										
All Races ^b	Both Sexes	7,194	5,578	5,671	5,046	1,392	430	23,872	25,504	+	+
	Males	6,655	5,284	5,439	4,842	1,369	396	22,579	24,134	+	+
	Females	539	294	232	204	23	34	1,293	1,370	+	+
White ^b	Both Sexes	4,740	4,115	4,237	4,257	1,088	369	17,633	18,977	+	+
	Males	4,353	3,896	4,073	4,087	1,088	346	16,692	17,970	+	+
	Females	387	219	164	170	0	23	941	1,007	+	+
Black ^b	Both Sexes	1,980	1,234	1,078	520	122	26	4,840	4,973	+	+
	Males	1,874	1,177	1,056	486	122	26	4,620	4,753	+	+
	Females	106	57	22	34	0	0	220	220	+	+
Asian/ Pacific Islander ^c	Both Sexes	118	97	79	+	+	+	339	+	+	+
	Males	118	91	79	+	+	+	333	+	+	+
	Females	0	6	0	+	+	+	6	+	+	+
Hispanic ^d	Both Sexes	1,227	1,113	1,058	+	+	+	4,149	+	+	+
	Males	1,093	1,071	1,017	+	+	+	3,916	+	+	+
	Females	134	42	41	+	+	+	233	+	+	+

Estimated prevalence percent^a on January 1, 2011, of the SEER^c population diagnosed in the previous 19 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)									Age-Adjusted ^f	
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>	<u>Sex</u>											
All Races ^c	Both Sexes	0.0084%	-	-	0.0012%	0.0058%	0.0205%	0.0188%	0.0111%	0.0093%	0.0145%	0.0081%
	Males	0.0162%	-	-	0.0021%	0.0114%	0.0403%	0.0381%	0.0224%	0.0177%	0.0275%	0.0160%
	Females	0.0007%	-	-	-	0.0002%	0.0008%	0.0004%	0.0009%	0.0026%	0.0069%	0.0006%
White ^c	Both Sexes	0.0085%	-	-	0.0008%	0.0049%	0.0203%	0.0194%	0.0117%	0.0100%	0.0153%	0.0080%
	Males	0.0163%	-	-	0.0014%	0.0093%	0.0390%	0.0387%	0.0232%	0.0189%	0.0288%	0.0156%
	Females	0.0008%	-	-	-	-	0.0009%	0.0003%	0.0009%	0.0026%	0.0073%	0.0007%
Black ^c	Both Sexes	0.0118%	-	-	0.0038%	0.0156%	0.0342%	0.0236%	0.0125%	0.0064%	0.0089%	0.0124%
	Males	0.0239%	-	-	0.0072%	0.0329%	0.0719%	0.0497%	0.0263%	0.0114%	-	0.0257%
	Females	0.0008%	-	-	-	-	-	-	-	-	-	0.0009%
Asian/ Pacific Islander ^c	Both Sexes	0.0021%	-	-	-	0.0011%	0.0047%	0.0050%	0.0023%	0.0041%	0.0044%	0.0020%
	Males	0.0043%	-	-	-	0.0024%	0.0097%	0.0107%	0.0051%	0.0094%	0.0092%	0.0044%
	Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	Both Sexes	0.0083%	-	-	0.0011%	0.0065%	0.0252%	0.0228%	0.0153%	0.0179%	0.0385%	0.0111%
	Males	0.0155%	-	-	0.0020%	0.0127%	0.0481%	0.0462%	0.0309%	0.0318%	0.0659%	0.0212%
	Females	0.0010%	-	-	-	-	0.0015%	-	0.0020%	0.0076%	0.0222%	0.0017%

^a US 2011 cancer prevalence counts are based on 2011 cancer prevalence proportions from the SEER registries and 1/1/2011 US population estimates based on the average of 2010 and 2011 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 36 years for 1975-2011 SEER 9 data; 19 years for 1992-2011 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 36 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <36 and >=36. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.