

Table 14.8
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1960-1963 ^a	-	-	-	-	-	-	-	-	-
1970-1973 ^a	-	-	-	-	-	-	-	-	-
1975-1977 ^b	3.4	1.8	6.3	3.4	1.9	5.8	1.8	0.0	5.6
1978-1980 ^b	3.3	2.2	5.4	3.3	2.2	4.8	3.6	3.0	5.0
1981-1983 ^b	3.6	2.9	5.0	3.8	3.0	5.2	2.8	2.1	4.7
1984-1986 ^b	5.7	3.7	9.5	5.7	3.7	9.1	4.4	2.1	9.1
1987-1989 ^b	4.8	3.2	7.9	5.6	3.4	9.2	2.7	1.6	4.9
1990-1992 ^b	5.6	4.9	7.0	6.2	5.1	8.3	2.6	2.0	3.6
1993-1995 ^b	5.5	5.3	6.0	5.2	5.0	5.6	4.0	2.8	7.3
1996-1998 ^b	8.5	8.3	9.1	8.4	7.6	10.2	4.6	5.9	2.0
1999-2003 ^b	12.7	12.5	13.0	11.8	11.7	12.1	8.5	7.4	11.3
2004-2010 ^b	17.7 ^f	17.6 ^f	17.7 ^f	17.1 ^f	17.4 ^f	16.4 ^f	12.9 ^f	11.6	16.5 ^f
5-Year Period Survival (Percent) ^{cd}									
2010	17.4	17.0	18.5	16.8	16.5	17.7	12.3	11.2	15.1
Stage Distribution (%) 2004-2010 ^{ee}									
All Stages									
Number of cases	37,857	27,478	10,379	26,048	19,020	7,028	4,699	3,476	1,223
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	42	42	42	42	42	41	42	42	42
Regional	27	28	24	26	27	24	28	28	27
Distant	18	18	18	18	18	18	21	21	19
Unstaged	13	12	16	14	13	17	10	9	13
5-Year Relative Survival (Percent), 2004-2010 ^c									
Age at Diagnosis:									
Ages <45	35.0	32.7	40.0	38.4	36.0	43.0	24.9	23.9	27.5 ^g
Ages 45-54	20.0	19.1	24.9	20.2	19.5	23.5	12.3	10.4	19.6
Ages 55-64	18.9	17.6	24.2	19.5	18.0	25.3	10.4	9.0	15.8
Ages 65-74	14.5	13.8	15.8	12.8	11.9	14.6	11.1	9.0	14.1
Ages 75+	6.4	6.7	6.0	5.0	5.0	4.9	4.8	4.7	4.9
Ages <65	20.8	19.4	26.8	21.3	19.9	27.4	12.5	10.8	19.0
Ages 65+	10.6	10.8	10.3	8.9	8.9	8.9	8.8	7.5	10.0
Stage ^e :									
All Stages	16.6	16.4	17.2	16.1	15.9	16.5	11.5	10.1	15.3
Localized	29.7	29.6	29.8	28.7	28.7	28.7	22.1	19.8	28.5
Regional	10.9	10.5	12.0	11.1	10.6	12.6	5.3	4.6	7.1
Distant	2.8	2.3	4.3	2.9	2.3	4.6	2.4	1.9	4.0
Unstaged	5.6	5.5	5.8	5.3	5.4	5.3	3.6	1.7	6.5

^a Based on End Results data from a series of hospital registries and one population-based registry.

^b SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Based on follow-up of patients into 2011.

^c SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Based on follow-up of patients into 2011.

^d Period survival provides a 2010 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2007-2009: 0-1 year survival), (2006-2008: 1-2 year survival), (2005-2007: 2-3 year survival), (2004-2006: 3-4 year survival), (2003-2005: 4-5 years survival).

^e Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.

^f The difference between 1975-1977 and 2004-2010 is statistically significant ($p < .05$).

^g The standard error is between 5 and 10 percentage points.

^h The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.