

Table 14.9
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	13.9	14.9	17.7	21.5	23.0	24.7	27.8	26.9	27.7	29.0	31.1	32.8	36.0	38.2	39.3	39.4	41.9	44.9	44.2	47.7
2-year	6.6	8.3	9.8	12.3	13.6	15.0	17.4	17.0	17.7	19.9	21.5	22.2	24.6	26.2	27.0	28.6	30.2	32.3	30.5	
3-year	4.8	6.0	7.2	8.7	9.5	10.8	12.4	13.1	14.3	15.4	15.8	17.3	20.3	20.7	21.3	23.2	23.8	25.4		
4-year	4.1	4.6	6.3	6.9	6.9	9.1	10.1	10.7	12.0	12.3	13.6	15.4	16.8	17.6	18.4	19.7	20.7			
5-year	3.4	3.8	5.3	5.5	5.7	8.1	8.1	9.4	10.1	11.6	11.8	13.9	15.3	14.6	16.8	16.8				
6-year	3.0	3.5	5.0	5.0	4.9	7.0	7.4	8.4	9.1	10.7	10.7	11.8	14.6	13.3	15.3					
7-year	2.8	3.1	4.7	4.5	4.4	6.6	6.6	6.9	8.3	9.9	9.6	11.2	12.9	12.6						
8-year	2.7	2.9	4.3	4.4	4.2	6.4	5.8	6.6	7.8	9.6	8.7	10.2	11.8							
9-year	2.6	2.8	4.2	4.2	4.1	5.9	5.6	5.6	7.5	9.3	8.3	9.9								
10-year	2.6	2.8	4.0	4.1	3.6	5.6	5.4	5.5	7.4	8.9	7.6									
11-year	2.6	2.6	3.9	3.9	3.2	5.5	5.4	4.8	6.9	8.6										
12-year	2.5	2.6	3.8	3.8	3.1	5.2	5.4	4.6	6.8											
13-year	2.3	2.4	3.8	3.7	3.1	5.0	5.4	4.5												
14-year	2.3	2.4	3.7	3.7	3.1	4.8	5.4													
15-year	2.2	2.4	3.7	3.6	3.0	3.9														
16-year	2.2	2.3	3.7	3.6	3.0															
17-year	2.1	2.3	3.7	3.5																
18-year	2.0	2.2	3.7																	
19-year	2.0	2.1	3.7																	
20-year	2.0	2.1	3.6																	

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).