

Table 14.10
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975– 1979	1980– 1984	1985– 1989	1990– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	11.4	12.9	14.8	20.1	21.3	24.8	26.8	25.6	28.8	28.3	30.6	32.1	35.0	37.4	40.1	39.8	41.0	45.5	45.0	48.8
2-year	4.8	6.5	7.4	11.3	12.1	15.9	16.5	16.1	17.6	19.2	20.3	21.1	24.6	25.5	27.7	29.6	29.5	32.7	31.2	
3-year	3.2	4.9	5.2	8.0	8.9	11.2	12.0	12.0	13.7	15.4	14.9	17.6	19.9	19.9	21.6	23.7	23.9	25.6		
4-year	2.7	3.4	4.5	6.3	6.2	9.2	9.6	10.6	11.4	12.2	12.6	15.5	16.7	17.4	18.5	19.9	20.2			
5-year	2.0	2.8	3.6	5.1	5.0	7.8	7.9	8.9	10.1	11.2	10.8	14.2	15.6	14.3	16.9	16.8				
6-year	1.7	2.6	3.3	4.7	4.2	6.8	7.3	7.6	8.9	10.4	9.9	11.9	14.8	12.9	15.4					
7-year	1.4	2.2	3.3	4.2	3.7	6.5	6.8	6.2	8.3	9.8	8.9	11.0	13.1	12.2						
8-year	1.4	1.9	3.0	4.1	3.5	6.3	5.5	6.1	7.7	9.7	7.7	9.8	11.8							
9-year	1.3	1.9	3.0	3.9	3.4	5.9	5.4	5.2	7.5	9.7	7.2	9.3								
10-year	1.3	1.9	2.9	3.7	2.8	5.4	5.4	5.2	7.4	9.3	6.8									
11-year	1.3	1.8	2.9	3.6	2.3	5.2	5.4	4.4	6.9	9.1										
12-year	1.3	1.7	2.7	3.5	2.1	4.9	5.4	4.3	6.9											
13-year	1.1	1.4	2.7	3.4	2.1	4.8	5.4	4.3												
14-year	1.1	1.4	2.6	3.4	2.1	4.4	5.3													
15-year	1.0	1.4	2.6	3.4	2.1	3.7														
16-year	1.0	1.4	2.6	3.4	2.1															
17-year	1.0	1.4	2.6	3.3																
18-year	1.0	1.4	2.6																	
19-year	1.0	1.4	2.6																	
20-year	1.0	1.4	2.6																	

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).