

Table 14.11
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	18.7	18.9	23.5	24.2	26.3	24.6	29.7	29.7	25.4	30.2	32.3	34.5	38.7	40.5	37.3	38.6	44.6	43.2	42.1	44.6
2-year	9.9	11.8	14.4	14.4	16.5	13.2	19.3	18.7	17.9	21.5	24.4	24.7	24.5	28.1	25.4	26.0	32.1	31.3	28.7	
3-year	7.8	8.2	11.3	10.1	10.4	9.9	13.3	15.2	15.3	15.3	17.7	16.7	21.1	22.9	20.6	21.9	23.5	24.6		
4-year	6.6	6.7	9.8	8.1	8.3	8.7	11.0	10.8	13.4	12.6	15.9	14.9	17.1	18.1	18.1	19.2	22.2			
5-year	6.1	5.6	8.7	6.3	7.0	8.5	8.4	10.2	10.4	12.3	14.0	13.4	14.6	15.3	16.6	16.9				
6-year	5.4	5.1	8.3	5.6	6.0	7.3	7.3	9.9	9.3	11.3	12.7	11.8	14.2	14.1	15.1					
7-year	5.4	4.7	7.6	5.1	5.7	6.7	6.2	8.4	8.2	10.0	11.3	11.6	12.2	13.7						
8-year	5.2	4.6	6.9	5.0	5.4	6.4	6.2	7.5	8.0	9.2	10.7	11.0	11.7							
9-year	4.9	4.5	6.5	5.0	5.4	5.9	5.8	6.2	7.4	8.3	10.6	11.0								
10-year	4.9	4.4	6.1	4.9	5.1	5.9	5.2	5.9	7.4	8.0	9.5									
11-year	4.9	4.3	5.9	4.4	4.8	5.9	5.2	5.6	6.9	7.5										
12-year	4.5	4.3	5.8	4.3	4.8	5.7	5.2	5.3	6.3											
13-year	4.4	4.2	5.7	4.3	4.8	5.4	5.2	5.0												
14-year	4.4	4.2	5.7	4.2	4.8	5.4	5.2													
15-year	4.4	4.2	5.7	3.9	4.4	4.3														
16-year	4.3	4.1	5.7	3.9	4.4															
17-year	4.0	3.9	5.7	3.7																
18-year	3.8	3.6	5.6																	
19-year	3.8	3.3	5.6																	
20-year	3.8	3.3	5.5																	

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).