


Table 14.18  
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.76	0.02		
High Five States				
D.C.	8.70 <sup>b</sup>	0.55	(01)	51.03 <sup>c</sup>
Louisiana	7.58 <sup>b</sup>	0.18	(02)	31.45 <sup>c</sup>
Texas	7.52 <sup>b</sup>	0.08	(03)	30.40 <sup>c</sup>
Hawaii	7.28 <sup>b</sup>	0.31	(04)	26.25 <sup>c</sup>
California	7.03 <sup>b</sup>	0.06	(05)	22.01 <sup>c</sup>
Low Five States				
South Dakota	4.27 <sup>b</sup>	0.30	(47)	-25.93 <sup>c</sup>
Utah	4.13 <sup>b</sup>	0.20	(48)	-28.37 <sup>c</sup>
Montana	4.07 <sup>b</sup>	0.26	(49)	-29.31 <sup>c</sup>
Idaho	4.01 <sup>b</sup>	0.23	(50)	-30.44 <sup>c</sup>
North Dakota	3.74 <sup>b</sup>	0.31	(51)	-35.15 <sup>c</sup>

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.94	0.15	(12)	2.98	Montana	4.07 <sup>b</sup>	0.26	(49)	-29.31 <sup>c</sup>
Alaska	5.71	0.49	(18)	-0.97	Nebraska	4.41 <sup>b</sup>	0.21	(46)	-23.52 <sup>c</sup>
Arizona	5.57	0.13	(22)	-3.40	Nevada	5.84	0.21	(14)	1.34
Arkansas	6.20	0.19	(08)	7.61	New Hampshire	4.53 <sup>b</sup>	0.25	(44)	-21.44 <sup>c</sup>
California	7.03 <sup>b</sup>	0.06	(05)	22.01 <sup>c</sup>	New Jersey	5.34 <sup>b</sup>	0.10	(26)	-7.39
Colorado	5.04 <sup>b</sup>	0.15	(35)	-12.51	New Mexico	6.45	0.24	(07)	11.86
Connecticut	5.04 <sup>b</sup>	0.16	(37)	-12.56	New York	5.78	0.07	(16)	0.29
Delaware	6.06	0.34	(11)	5.13	North Carolina	5.22 <sup>b</sup>	0.10	(31)	-9.38
D.C.	8.70 <sup>b</sup>	0.55	(01)	51.03 <sup>c</sup>	North Dakota	3.74 <sup>b</sup>	0.31	(51)	-35.15 <sup>c</sup>
Florida	5.53 <sup>b</sup>	0.07	(24)	-4.04	Ohio	5.06 <sup>b</sup>	0.09	(34)	-12.12
Georgia	5.22 <sup>b</sup>	0.11	(32)	-9.49	Oklahoma	5.69	0.17	(19)	-1.24
Hawaii	7.28 <sup>b</sup>	0.31	(04)	26.25 <sup>c</sup>	Oregon	5.62	0.16	(20)	-2.45
Idaho	4.01 <sup>b</sup>	0.23	(50)	-30.44 <sup>c</sup>	Pennsylvania	5.32 <sup>b</sup>	0.08	(27)	-7.65
Illinois	5.46	0.09	(25)	-5.18	Rhode Island	5.92	0.31	(13)	2.67
Indiana	4.74 <sup>b</sup>	0.12	(40)	-17.75 <sup>c</sup>	South Carolina	5.61	0.15	(21)	-2.65
Iowa	4.55 <sup>b</sup>	0.16	(43)	-20.99 <sup>c</sup>	South Dakota	4.27 <sup>b</sup>	0.30	(47)	-25.93 <sup>c</sup>
Kansas	4.51 <sup>b</sup>	0.17	(45)	-21.73 <sup>c</sup>	Tennessee	5.84	0.13	(15)	1.34
Kentucky	5.22 <sup>b</sup>	0.15	(33)	-9.49	Texas	7.52 <sup>b</sup>	0.08	(03)	30.40 <sup>c</sup>
Louisiana	7.58 <sup>b</sup>	0.18	(02)	31.45 <sup>c</sup>	Utah	4.13 <sup>b</sup>	0.20	(48)	-28.37 <sup>c</sup>
Maine	4.70 <sup>b</sup>	0.24	(41)	-18.46 <sup>c</sup>	Vermont	5.31	0.39	(28)	-7.79
Maryland	5.76	0.14	(17)	0.02	Virginia	5.31 <sup>b</sup>	0.11	(29)	-7.95
Massachusetts	6.13	0.13	(10)	6.34	Washington	6.17	0.13	(09)	7.08
Michigan	5.30 <sup>b</sup>	0.10	(30)	-8.11	West Virginia	4.92 <sup>b</sup>	0.21	(39)	-14.71
Minnesota	4.93 <sup>b</sup>	0.13	(38)	-14.46	Wisconsin	5.04 <sup>b</sup>	0.13	(36)	-12.56
Mississippi	6.84 <sup>b</sup>	0.21	(06)	18.61 <sup>c</sup>	Wyoming	4.61	0.40	(42)	-19.93 <sup>c</sup>
Missouri	5.54	0.13	(23)	-3.86					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant (p<=.0002).  
<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.  
SE Standard error of the rate.  
PD Percent difference between state rate and total U.S. rate.  
- Statistic not shown. Rate based on less than 16 cases for the time interval.