


Table 14.19
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	8.54	0.03		
High Five States				
D.C.	13.97 ^b	1.05	(01)	63.50 ^c
Louisiana	11.57 ^b	0.34	(02)	35.42 ^c
Texas	11.00 ^b	0.15	(03)	28.73 ^c
Hawaii	10.47 ^b	0.54	(04)	22.54 ^c
Mississippi	10.27 ^b	0.40	(05)	20.24 ^c
Low Five States				
Utah	5.68 ^b	0.34	(47)	-33.56 ^c
Idaho	5.53 ^b	0.39	(48)	-35.27 ^c
Montana	5.47 ^b	0.44	(49)	-35.92 ^c
North Dakota	5.43 ^b	0.55	(50)	-36.49 ^c
South Dakota	5.19 ^b	0.49	(51)	-39.24 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	8.86	0.28	(14)	3.74	Montana	5.47 ^b	0.44	(49)	-35.92 ^c
Alaska	7.97	0.86	(29)	-6.73	Nebraska	6.49 ^b	0.38	(45)	-24.06 ^c
Arizona	7.81	0.22	(30)	-8.60	Nevada	8.40	0.38	(18)	-1.64
Arkansas	9.20	0.35	(10)	7.69	New Hampshire	6.59 ^b	0.45	(42)	-22.81 ^c
California	10.26 ^b	0.11	(06)	20.08 ^c	New Jersey	7.99	0.20	(28)	-6.51
Colorado	7.40 ^b	0.27	(36)	-13.41	New Mexico	9.34	0.43	(09)	9.34
Connecticut	7.63	0.29	(35)	-10.67	New York	8.75	0.14	(16)	2.39
Delaware	9.50	0.64	(08)	11.23	North Carolina	8.00	0.19	(27)	-6.39
D.C.	13.97 ^b	1.05	(01)	63.50 ^c	North Dakota	5.43 ^b	0.55	(50)	-36.49 ^c
Florida	8.37	0.12	(19)	-1.98	Ohio	7.65 ^b	0.16	(34)	-10.47
Georgia	7.70 ^b	0.21	(32)	-9.82	Oklahoma	8.21	0.30	(21)	-3.87
Hawaii	10.47 ^b	0.54	(04)	22.54 ^c	Oregon	8.16	0.28	(22)	-4.49
Idaho	5.53 ^b	0.39	(48)	-35.27 ^c	Pennsylvania	8.00 ^b	0.15	(26)	-6.32
Illinois	8.07	0.17	(25)	-5.53	Rhode Island	8.88	0.57	(13)	3.92
Indiana	6.89 ^b	0.21	(40)	-19.35 ^c	South Carolina	8.47	0.28	(17)	-0.80
Iowa	6.35 ^b	0.28	(46)	-25.62 ^c	South Dakota	5.19 ^b	0.49	(51)	-39.24 ^c
Kansas	6.56 ^b	0.30	(43)	-23.23 ^c	Tennessee	8.99	0.25	(11)	5.27
Kentucky	7.78	0.28	(31)	-8.94	Texas	11.00 ^b	0.15	(03)	28.73 ^c
Louisiana	11.57 ^b	0.34	(02)	35.42 ^c	Utah	5.68 ^b	0.34	(47)	-33.56 ^c
Maine	6.94 ^b	0.43	(39)	-18.81 ^c	Vermont	8.34	0.72	(20)	-2.39
Maryland	8.83	0.26	(15)	3.34	Virginia	8.12	0.22	(24)	-5.00
Massachusetts	9.73 ^b	0.25	(07)	13.86	Washington	8.99	0.24	(12)	5.21
Michigan	7.70 ^b	0.18	(33)	-9.83	West Virginia	6.88 ^b	0.37	(41)	-19.49 ^c
Minnesota	7.16 ^b	0.24	(38)	-16.22 ^c	Wisconsin	7.35 ^b	0.23	(37)	-14.02
Mississippi	10.27 ^b	0.40	(05)	20.24 ^c	Wyoming	6.51	0.71	(44)	-23.75 ^c
Missouri	8.16	0.23	(23)	-4.54					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.