

Table 17.7
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	40.8	43.7	42.3	48.0	38.8	32.3	46.3	54.5	41.7	37.1	52.3	35.7	51.3	46.1	43.6	56.3	46.0	55.3	42.9	42.5
2-year	31.1	28.3	24.5	31.2	23.1	26.8	27.9	34.8	36.9	19.0	34.9	19.4	32.9	22.3	31.2	44.8	26.6	21.6	27.2	
3-year	24.4	19.4	18.9	24.4	21.2	21.3	17.2	26.0	34.9	16.8	22.9	9.9	24.8	14.3	27.1	36.7	12.6	12.5		
4-year	21.5	14.4	15.4	21.3	19.3	11.8	14.4	21.6	32.8	8.6	18.9	9.9	16.0	14.3	22.7	27.9	8.4			
5-year	20.1	13.3	13.4	20.2	17.6	9.8	14.4	21.6	22.8	0.0	18.9	7.8	14.2	12.4	18.5	24.5				
6-year	17.1	10.5	11.9	16.9	15.2	7.9	14.4	21.6	20.2	0.0	17.0	7.8	12.3	12.4	18.5					
7-year	16.3	10.5	10.8	16.5	15.2	6.0	8.9	20.6	17.6	0.0	15.0	7.8	10.3	10.4						
8-year	15.7	9.9	8.7	15.8	15.2	6.0	8.9	20.6	17.6	0.0	15.0	7.8	5.2							
9-year	15.2	9.5	8.7	15.3	12.9	4.0	8.9	20.6	15.0	0.0	15.0	7.8								
10-year	14.4	7.7	8.4	14.0	12.9	4.0	8.9	20.2	15.0	0.0	11.2									
11-year	13.7	7.7	8.4	13.7	12.9	4.0	8.9	18.1	15.0	0.0										
12-year	13.7	7.7	8.1	12.7	10.6	4.0	8.9	12.6	15.0											
13-year	13.7	7.7	8.1	11.8	10.6	4.0	8.9	12.6												
14-year	13.7	7.7	8.1	11.8	10.6	4.0	8.9													
15-year	13.7	6.8	8.1	11.6	10.6	4.0														
16-year	12.9	6.8	8.1	11.6	10.6															
17-year	12.3	6.8	8.1	11.6																
18-year	12.3	6.8	7.9																	
19-year	12.3	6.8	7.9																	
20-year	12.3	6.5	7.4																	

^a

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).