

Table 17.9
Mesothelioma (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.21	0.17	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.10	0.07
	10	0.00	0.00	0.00	0.21	0.17		10	0.00	0.00	0.00	0.10	0.07
	20	0.00	0.00	0.00	0.21	0.17		20	0.00	0.00	0.00	0.10	0.07
	30	0.00	0.00	0.01	0.21	0.18		30	0.00	0.00	0.00	0.10	0.07
	40	0.00	0.01	0.05	0.22	0.18		40	0.00	0.00	0.02	0.10	0.07
	50	0.01	0.05	0.13	0.22	0.18		50	0.00	0.02	0.05	0.10	0.07
	60	0.04	0.13	0.21	0.23	0.19		60	0.02	0.05	0.09	0.10	0.07
	70	0.10	0.19	-	0.21	0.18		70	0.04	0.08	-	0.10	0.07
White	80	0.14	-	-	0.17	0.14	American Indian/ Alaska Native ^a	80	0.05	-	-	0.07	0.05
	0	0.00	0.00	0.00	0.23	0.19		0	0.00	0.00	0.00	0.08	0.14
	10	0.00	0.00	0.00	0.23	0.19		10	0.00	0.00	0.00	0.09	0.14
	20	0.00	0.00	0.00	0.23	0.19		20	0.00	0.00	0.00	0.09	0.14
	30	0.00	0.00	0.02	0.24	0.19		30	0.00	0.00	0.01	0.09	0.14
	40	0.00	0.02	0.06	0.24	0.19		40	0.00	0.01	0.01	0.09	0.15
	50	0.01	0.06	0.14	0.24	0.20		50	0.01	0.01	0.01	0.09	0.15
	60	0.05	0.14	0.23	0.25	0.21		60	0.00	0.01	0.09	0.09	0.17
Black	70	0.11	0.21	-	0.23	0.20	Hispanic ^b	70	0.01	0.10	-	0.11	0.18
	80	0.15	-	-	0.18	0.15		80	0.14	-	-	0.15	0.18
	0	0.00	0.00	0.00	0.10	0.06		0	0.00	0.00	0.00	0.19	0.10
	10	0.00	0.00	0.00	0.10	0.06		10	0.00	0.00	0.00	0.19	0.10
	20	0.00	0.00	0.00	0.10	0.06		20	0.00	0.00	0.00	0.19	0.10
	30	0.00	0.00	0.01	0.10	0.06		30	0.00	0.00	0.02	0.19	0.10
	40	0.00	0.01	0.03	0.11	0.07		40	0.00	0.01	0.05	0.19	0.10
	50	0.01	0.03	0.07	0.11	0.07		50	0.01	0.05	0.12	0.20	0.11
	60	0.02	0.07	0.11	0.11	0.07		60	0.04	0.12	0.19	0.19	0.11
	70	0.06	0.11	-	0.12	0.07		70	0.09	0.17	-	0.18	0.10
	80	0.07	-	-	0.09	0.05		80	0.11	-	-	0.12	0.07

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.