

Table 17.10
Mesothelioma (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Females, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.05	0.04	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.02	0.01
	10	0.00	0.00	0.00	0.05	0.04		10	0.00	0.00	0.00	0.02	0.01
	20	0.00	0.00	0.00	0.05	0.04		20	0.00	0.00	0.00	0.02	0.01
	30	0.00	0.00	0.01	0.05	0.04		30	0.00	0.00	0.00	0.02	0.01
	40	0.00	0.01	0.02	0.05	0.04		40	0.00	0.00	0.00	0.02	0.01
	50	0.00	0.01	0.03	0.05	0.04		50	0.00	0.00	0.01	0.02	0.01
	60	0.01	0.03	0.05	0.05	0.04		60	0.00	0.01	0.02	0.02	0.01
	70	0.02	0.04	-	0.04	0.03		70	0.01	0.01	-	0.02	0.01
White	80	0.02	-	-	0.03	0.02	American Indian/ Alaska Native ^a	80	0.01	-	-	0.01	0.01
	0	0.00	0.00	0.00	0.06	0.04		0	0.00	0.00	0.00	0.04	0.02
	10	0.00	0.00	0.00	0.06	0.04		10	0.00	0.00	0.00	0.04	0.02
	20	0.00	0.00	0.00	0.06	0.04		20	0.00	0.00	0.00	0.04	0.02
	30	0.00	0.00	0.01	0.06	0.04		30	0.00	0.00	0.00	0.04	0.02
	40	0.00	0.01	0.02	0.06	0.04		40	0.00	0.00	0.00	0.04	0.02
	50	0.00	0.02	0.04	0.06	0.04		50	0.00	0.00	0.00	0.04	0.03
	60	0.01	0.03	0.05	0.05	0.04		60	0.00	0.00	0.04	0.04	0.03
Black	70	0.02	0.04	-	0.05	0.04	Hispanic ^b	70	0.00	0.04	-	0.05	0.03
	80	0.03	-	-	0.03	0.03		80	0.05	-	-	0.06	0.02
	0	0.00	0.00	0.00	0.03	0.01		0	0.00	0.00	0.00	0.05	0.02
	10	0.00	0.00	0.00	0.03	0.01		10	0.00	0.00	0.00	0.05	0.02
	20	0.00	0.00	0.00	0.03	0.01		20	0.00	0.00	0.00	0.05	0.02
	30	0.00	0.00	0.01	0.03	0.01		30	0.00	0.00	0.01	0.05	0.02
	40	0.00	0.01	0.01	0.03	0.01		40	0.00	0.01	0.02	0.05	0.02
	50	0.00	0.01	0.02	0.02	0.01		50	0.00	0.01	0.03	0.05	0.02
	60	0.00	0.01	0.02	0.02	0.01		60	0.01	0.03	0.04	0.04	0.02
	70	0.01	0.02	-	0.02	0.01		70	0.02	0.03	-	0.04	0.02
	80	0.02	-	-	0.02	0.01		80	0.02	-	-	0.02	0.01

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.