

Table 18.9
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	68.0	69.5	72.3	71.9	73.9	70.6	74.3	74.8	73.5	72.4	72.5	73.4	77.1	74.5	75.8	73.8	76.6	77.1	81.4	80.7
2-year	52.1	54.5	57.6	57.4	58.8	57.6	60.5	62.2	59.6	59.9	59.5	62.7	64.9	66.0	64.3	65.9	65.9	67.7	71.5	
3-year	39.6	42.9	45.2	46.2	47.9	44.9	48.6	50.9	47.7	50.0	49.2	53.2	55.7	58.1	56.7	57.6	56.8	60.0		
4-year	31.5	34.2	35.7	36.9	40.1	38.2	38.7	39.6	38.6	41.3	43.7	46.5	47.6	50.2	49.8	50.4	49.1			
5-year	25.2	26.9	27.4	29.8	33.4	31.8	31.3	33.9	31.0	34.4	37.1	39.5	41.9	44.7	45.5	45.1				
6-year	20.1	21.0	22.1	24.5	28.3	27.1	26.4	29.5	26.8	29.2	32.3	34.8	37.3	39.1	40.4					
7-year	16.2	16.7	17.5	20.4	23.7	23.7	22.5	24.1	23.1	25.3	28.8	31.1	32.8	34.5						
8-year	13.6	13.4	14.9	17.3	20.4	21.4	19.8	21.9	19.7	22.6	25.2	27.9	29.5							
9-year	11.5	11.2	11.9	15.3	18.3	19.5	17.1	19.1	17.9	20.4	22.6	24.9								
10-year	9.9	9.7	10.4	13.6	17.2	17.2	15.5	17.5	15.9	19.3	20.6									
11-year	8.8	8.7	9.1	12.6	15.6	15.6	14.3	16.3	13.6	18.3										
12-year	7.8	7.7	8.3	11.0	14.3	14.9	13.0	14.4	13.1											
13-year	6.7	6.6	7.6	10.2	13.6	13.3	11.8	13.5												
14-year	6.1	6.1	6.9	9.4	12.3	12.6	11.1													
15-year	5.3	5.7	6.1	9.1	11.9	11.5														
16-year	4.9	5.2	5.7	8.5	10.7															
17-year	4.6	4.7	5.4	8.0																
18-year	4.1	4.3	5.1																	
19-year	3.9	3.8	4.9																	
20-year	3.4	3.5	4.7																	

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).