

Table 18.10  
MyelomaSEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	67.7	69.6	73.2	72.5	74.2	71.2	74.0	76.1	75.5	73.6	74.1	74.1	76.9	76.5	75.9	73.5	77.0	76.1	83.2	82.7
2-year	52.2	54.6	59.3	58.4	59.8	58.3	60.9	63.4	61.2	61.6	61.4	61.4	65.3	67.8	63.8	66.6	67.3	66.7	72.8	
3-year	38.9	42.8	46.2	46.6	49.2	45.0	50.2	53.3	50.5	52.6	50.7	52.3	56.4	60.0	56.0	57.9	58.9	57.8		
4-year	30.9	33.6	36.6	36.9	41.2	39.8	40.7	43.4	42.6	43.3	45.7	46.5	47.8	51.2	49.7	51.3	50.6			
5-year	24.3	25.8	28.4	30.2	36.2	33.7	31.5	37.8	34.4	35.8	39.5	41.3	42.2	45.6	46.1	46.7				
6-year	19.3	20.8	23.5	25.1	31.6	29.7	26.4	33.6	29.6	30.0	34.6	36.0	37.7	39.7	40.3					
7-year	15.4	16.9	19.1	21.2	26.0	25.8	22.3	27.7	25.7	26.1	30.8	32.5	34.2	35.2						
8-year	13.1	13.8	16.2	18.5	22.0	24.0	20.1	25.2	21.3	23.3	26.8	29.2	30.5							
9-year	11.1	11.7	12.9	16.6	20.0	21.2	18.1	22.9	19.0	19.7	24.2	26.0								
10-year	9.3	9.9	11.2	14.7	19.0	18.3	16.4	21.2	17.0	18.5	21.6									
11-year	7.6	8.9	9.9	13.5	17.4	17.0	14.9	20.3	13.9	17.0										
12-year	7.0	7.7	9.4	11.9	15.5	16.6	14.3	18.2	13.1											
13-year	5.9	6.7	8.6	11.1	14.8	14.5	13.1	16.6												
14-year	5.3	6.1	7.9	10.3	13.1	14.1	12.6													
15-year	4.7	5.7	7.1	9.8	12.9	12.6														
16-year	4.5	5.2	6.7	9.5	11.2															
17-year	4.0	4.8	6.4	8.7																
18-year	3.6	4.4	6.2																	
19-year	3.4	4.2	6.2																	
20-year	2.7	3.9	6.0																	

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).