

Table 18.11
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	68.2	69.3	71.2	71.1	73.6	69.9	74.6	73.3	71.1	71.0	70.7	72.6	77.4	72.1	75.6	74.2	76.1	78.1	79.2	78.2
2-year	52.0	54.3	55.9	56.3	57.8	56.8	60.1	60.7	57.9	57.9	57.5	64.3	64.4	64.0	64.9	65.1	64.0	68.8	70.0	
3-year	40.4	42.9	44.3	45.7	46.3	44.9	46.9	48.2	44.6	47.1	47.6	54.4	54.8	56.0	57.6	57.2	53.9	62.4		
4-year	32.2	34.9	34.7	36.8	39.0	36.4	36.5	35.3	34.1	38.9	41.5	46.5	47.3	49.0	50.0	49.4	47.0			
5-year	26.1	28.1	26.5	29.4	30.2	29.8	31.1	29.6	27.2	32.8	34.4	37.3	41.3	43.5	44.8	43.4				
6-year	21.0	21.2	20.7	23.8	24.7	24.4	26.3	24.8	23.7	28.2	29.7	33.3	36.9	38.2	40.4					
7-year	17.1	16.5	15.8	19.4	21.3	21.5	22.7	20.1	20.1	24.4	26.4	29.4	31.2	33.7						
8-year	14.1	13.0	13.6	15.9	18.5	18.6	19.5	18.2	17.9	21.7	23.4	26.3	28.3							
9-year	12.0	10.8	10.8	13.9	16.4	17.7	16.1	15.0	16.5	21.2	20.9	23.5								
10-year	10.5	9.4	9.5	12.4	15.1	16.0	14.6	13.3	14.6	20.1	19.4									
11-year	10.0	8.4	8.4	11.6	13.7	14.2	13.6	11.8	13.2	19.6										
12-year	8.7	7.6	7.3	10.1	12.8	13.1	11.4	10.4	13.0											
13-year	7.5	6.6	6.6	9.2	12.0	12.0	10.3	10.1												
14-year	6.9	6.0	5.9	8.4	11.4	11.0	9.5													
15-year	6.0	5.7	5.2	8.2	10.8	10.2														
16-year	5.4	5.1	4.6	7.4	10.1															
17-year	5.2	4.6	4.4	7.1																
18-year	4.5	4.2	4.0																	
19-year	4.3	3.4	3.7																	
20-year	3.9	3.1	3.4																	

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).