

Table 18.12
Myeloma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Both Sexes, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.72	0.42	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.54	0.29
	10	0.00	0.00	0.01	0.73	0.42		10	0.00	0.00	0.00	0.54	0.30
	20	0.00	0.01	0.04	0.73	0.42		20	0.00	0.00	0.02	0.54	0.30
	30	0.01	0.03	0.11	0.73	0.43		30	0.00	0.02	0.07	0.54	0.30
	40	0.03	0.11	0.28	0.74	0.43		40	0.02	0.06	0.16	0.54	0.30
	50	0.08	0.26	0.51	0.73	0.44		50	0.05	0.15	0.33	0.54	0.30
	60	0.18	0.45	0.64	0.68	0.44		60	0.10	0.29	0.44	0.50	0.29
	70	0.31	0.52	-	0.56	0.40		70	0.20	0.37	-	0.43	0.27
80	0.29	-	-	0.35	0.31	80	0.20	-	-	0.27	0.20		
White	0	0.00	0.00	0.00	0.66	0.39	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.44	0.31
	10	0.00	0.00	0.01	0.67	0.40		10	0.00	0.00	0.01	0.45	0.32
	20	0.00	0.01	0.03	0.67	0.40		20	0.00	0.01	0.03	0.45	0.32
	30	0.01	0.03	0.10	0.67	0.40		30	0.01	0.03	0.06	0.46	0.33
	40	0.02	0.09	0.25	0.68	0.41		40	0.02	0.06	0.16	0.46	0.33
	50	0.07	0.23	0.47	0.67	0.41		50	0.04	0.14	0.30	0.46	0.35
	60	0.17	0.42	0.60	0.63	0.41		60	0.11	0.29	0.39	0.45	0.35
	70	0.28	0.49	-	0.53	0.38		70	0.20	0.31	-	0.38	0.32
80	0.27	-	-	0.33	0.29	80	0.16	-	-	0.25	0.24		
Black	0	0.00	0.00	0.00	1.21	0.66	Hispanic ^b	0	0.00	0.00	0.00	0.76	0.42
	10	0.00	0.00	0.01	1.24	0.67		10	0.00	0.00	0.01	0.77	0.43
	20	0.00	0.01	0.07	1.24	0.67		20	0.00	0.01	0.03	0.77	0.43
	30	0.01	0.07	0.23	1.26	0.68		30	0.01	0.03	0.11	0.77	0.43
	40	0.06	0.23	0.56	1.27	0.69		40	0.03	0.10	0.26	0.77	0.43
	50	0.17	0.52	0.95	1.26	0.71		50	0.08	0.24	0.49	0.76	0.44
	60	0.39	0.85	1.14	1.20	0.71		60	0.17	0.43	0.64	0.72	0.43
	70	0.56	0.91	-	0.98	0.66		70	0.29	0.52	-	0.60	0.39
80	0.51	-	-	0.61	0.51	80	0.30	-	-	0.40	0.29		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.