

Table 18.13
Myeloma

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2009-2011 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.00	0.00	0.83	0.47	Asian/ Pacific Islander	0	0.00	0.00	0.00	0.63	0.36
	10	0.00	0.00	0.01	0.84	0.47		10	0.00	0.00	0.00	0.63	0.36
	20	0.00	0.01	0.04	0.84	0.48		20	0.00	0.00	0.02	0.63	0.36
	30	0.01	0.04	0.13	0.85	0.48		30	0.00	0.02	0.08	0.64	0.36
	40	0.03	0.12	0.32	0.86	0.49		40	0.02	0.07	0.19	0.64	0.37
	50	0.09	0.30	0.61	0.85	0.50		50	0.06	0.17	0.40	0.63	0.37
	60	0.22	0.55	0.78	0.82	0.50		60	0.12	0.36	0.55	0.59	0.36
	70	0.39	0.64	-	0.69	0.47		70	0.26	0.47	-	0.52	0.34
80	0.37	-	-	0.44	0.37	80	0.26	-	-	0.32	0.26		
White	0	0.00	0.00	0.00	0.78	0.45	American Indian/ Alaska Native ^a	0	0.00	0.00	0.00	0.48	0.31
	10	0.00	0.00	0.01	0.79	0.45		10	0.00	0.00	0.01	0.49	0.31
	20	0.00	0.01	0.04	0.79	0.45		20	0.00	0.01	0.04	0.49	0.31
	30	0.01	0.03	0.11	0.80	0.46		30	0.01	0.04	0.07	0.50	0.32
	40	0.03	0.11	0.29	0.81	0.47		40	0.04	0.07	0.18	0.51	0.33
	50	0.08	0.27	0.57	0.80	0.47		50	0.04	0.15	0.32	0.50	0.34
	60	0.20	0.52	0.73	0.77	0.48		60	0.12	0.30	0.43	0.50	0.36
	70	0.36	0.61	-	0.66	0.46		70	0.21	0.36	-	0.44	0.34
80	0.36	-	-	0.43	0.36	80	0.21	-	-	0.34	0.24		
Black	0	0.00	0.00	0.00	1.25	0.68	Hispanic ^b	0	0.00	0.00	0.00	0.88	0.48
	10	0.00	0.00	0.01	1.28	0.70		10	0.00	0.00	0.01	0.89	0.49
	20	0.00	0.01	0.07	1.29	0.71		20	0.00	0.01	0.04	0.89	0.49
	30	0.01	0.07	0.24	1.31	0.72		30	0.01	0.04	0.12	0.89	0.50
	40	0.06	0.24	0.62	1.33	0.74		40	0.03	0.11	0.30	0.90	0.50
	50	0.18	0.58	1.05	1.33	0.75		50	0.08	0.27	0.57	0.89	0.51
	60	0.45	0.98	1.25	1.29	0.77		60	0.20	0.52	0.75	0.85	0.51
	70	0.67	1.02	-	1.08	0.74		70	0.36	0.62	-	0.74	0.47
80	0.57	-	-	0.65	0.59	80	0.36	-	-	0.51	0.37		

Devcan 6.8.0, August 2014, National Cancer Institute (<http://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, and Georgia excluding ATL/RG). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.