


Table 18.18
Myeloma

Age-Adjusted Cancer Death^a Rates By State, All Races, 2007-2011

Males and Females

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
TOTAL U.S.	3.36	0.01							
High Five States									
D.C.	5.45 ^b	0.45	(01)	62.30 ^c					
Alabama	3.97 ^b	0.12	(02)	18.29 ^c					
Louisiana	3.83 ^b	0.13	(03)	14.08					
North Carolina	3.83 ^b	0.09	(04)	13.94					
Maryland	3.80 ^b	0.12	(05)	13.18					
Low Five States									
Rhode Island	2.86	0.22	(47)	-14.82					
Connecticut	2.86 ^b	0.12	(48)	-14.83					
New Mexico	2.84	0.16	(49)	-15.46 ^c					
Nevada	2.77 ^b	0.15	(50)	-17.68 ^c					
Hawaii	2.46 ^b	0.18	(51)	-26.71 ^c					

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.97 ^b	0.12	(02)	18.29 ^c	Montana	3.26	0.24	(35)	-2.89
Alaska	2.91	0.39	(46)	-13.26	Nebraska	3.32	0.18	(32)	-1.30
Arizona	2.96 ^b	0.09	(45)	-11.90	Nevada	2.77 ^b	0.15	(50)	-17.68 ^c
Arkansas	3.50	0.15	(22)	4.30	New Hampshire	3.34	0.22	(28)	-0.57
California	3.14 ^b	0.04	(39)	-6.58	New Jersey	3.22	0.08	(38)	-4.05
Colorado	3.09	0.12	(41)	-8.12	New Mexico	2.84	0.16	(49)	-15.46 ^c
Connecticut	2.86 ^b	0.12	(48)	-14.83	New York	3.11 ^b	0.05	(40)	-7.28
Delaware	3.71	0.27	(09)	10.39	North Carolina	3.83 ^b	0.09	(04)	13.94
D.C.	5.45 ^b	0.45	(01)	62.30 ^c	North Dakota	3.58	0.30	(13)	6.51
Florida	2.98 ^b	0.05	(44)	-11.40	Ohio	3.55	0.07	(16)	5.79
Georgia	3.57	0.09	(15)	6.25	Oklahoma	3.53	0.13	(19)	5.20
Hawaii	2.46 ^b	0.18	(51)	-26.71 ^c	Oregon	3.55	0.13	(18)	5.59
Idaho	3.27	0.21	(33)	-2.54	Pennsylvania	3.36	0.07	(27)	0.01
Illinois	3.41	0.07	(26)	1.55	Rhode Island	2.86	0.22	(47)	-14.82
Indiana	3.53	0.10	(20)	5.19	South Carolina	3.75	0.13	(08)	11.78
Iowa	3.32	0.14	(30)	-1.18	South Dakota	3.33	0.27	(29)	-0.95
Kansas	3.66	0.16	(11)	8.91	Tennessee	3.78 ^b	0.11	(07)	12.49
Kentucky	3.42	0.12	(24)	1.79	Texas	3.23	0.06	(37)	-3.82
Louisiana	3.83 ^b	0.13	(03)	14.08	Utah	3.42	0.19	(25)	1.79
Maine	3.06	0.19	(42)	-9.02	Vermont	2.98	0.29	(43)	-11.35
Maryland	3.80 ^b	0.12	(05)	13.18	Virginia	3.80 ^b	0.10	(06)	13.07
Massachusetts	3.25	0.09	(36)	-3.28	Washington	3.32	0.10	(31)	-1.29
Michigan	3.70 ^b	0.08	(10)	10.13	West Virginia	3.51	0.17	(21)	4.58
Minnesota	3.49	0.11	(23)	4.05	Wisconsin	3.64	0.11	(12)	8.32
Mississippi	3.55	0.15	(17)	5.59	Wyoming	3.27	0.35	(34)	-2.74
Missouri	3.58	0.10	(14)	6.48					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.