

Table 19.10  
Non-Hodgkin Lymphoma

SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	69.4	72.7	68.9	67.1	66.6	71.1	74.7	74.9	75.8	76.4	77.6	80.6	79.6	80.1	80.9	80.9	81.3	83.4	82.5	83.4
2-year	59.4	63.7	60.4	58.2	58.3	63.5	67.2	68.2	69.1	70.1	71.2	75.2	74.9	75.4	76.0	76.4	77.6	77.6	77.7	
3-year	53.3	58.0	55.7	53.7	54.5	59.7	63.8	64.5	64.3	66.6	67.9	71.9	72.5	72.5	72.9	73.5	75.2	74.8		
4-year	49.3	53.9	51.5	50.4	51.5	56.1	61.0	62.0	61.5	63.7	65.6	70.0	70.8	70.0	70.8	71.2	73.1			
5-year	45.4	50.1	48.2	47.6	48.5	53.6	58.2	59.2	59.0	61.5	63.5	67.9	68.9	68.7	69.6	68.8				
6-year	42.5	46.6	45.4	45.5	46.8	51.0	56.1	56.5	56.2	60.0	62.0	66.7	66.8	67.4	68.1					
7-year	39.9	44.2	43.1	43.6	44.8	49.1	54.2	54.5	54.6	58.5	60.0	65.2	65.6	65.3						
8-year	37.7	42.0	40.9	41.7	43.2	47.4	52.0	53.4	53.5	56.4	58.1	63.6	64.3							
9-year	35.6	40.1	39.0	40.2	42.3	46.5	50.4	51.5	52.3	55.0	57.1	62.7								
10-year	33.9	38.0	37.6	39.1	41.5	45.5	48.9	50.4	51.5	53.9	55.8									
11-year	32.7	36.3	36.3	38.0	40.7	43.7	47.6	49.5	50.7	52.4										
12-year	31.8	35.2	35.0	37.2	39.4	42.6	47.1	48.2	49.7											
13-year	30.6	34.2	34.1	36.1	38.8	41.8	45.5	46.9												
14-year	29.4	33.0	33.1	34.9	37.7	41.3	44.2													
15-year	28.0	31.9	32.3	34.3	36.7	40.8														
16-year	27.0	30.7	31.5	33.2	35.9															
17-year	26.4	29.9	30.7	32.5																
18-year	26.0	29.3	30.1																	
19-year	25.4	28.7	29.3																	
20-year	24.9	27.9	29.0																	

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).